

Virtual Reality as Distraction Technique for Pain Management in Children and Adolescents

M**Barbara Atzori***University of Florence, Italy***Hunter G. Hoffman***University of Washington, USA***Laura Vagnoli***Meyer Children's Hospital of Florence, Italy***Andrea Messeri***Meyer Children's Hospital of Florence, Italy***Rosapia Lauro Grotto***University of Florence, Italy*

INTRODUCTION

The problem: uncontrolled pain during medical procedures. For many medical procedures, patients remain awake during the procedure, they feel pain during the procedure, and they remember pain and unpleasantness they experienced, long after the procedure is over. These memories for the painful experience(s) can influence the patient's attitudes towards receiving healthcare, increases avoidance of medical visits that could help prevent medical problems. For example, a child who has a very painful or anxiety provoking dental procedure is more likely to become avoidant of dentists. The consequences of avoidance can be expensive. Many dental problems that could be easily prevented or corrected early on with regular checkups (e.g, filling a cavity), become much more expensive and painful problems, as the disease progresses untreated (e.g., root canal and tooth extraction) for patients who refuse to visit the dentist. Excessive pain during medical procedures is a worldwide problem, and affects a wide range of patient populations. For example, patients

with a wide range of medical problems often get intravenous (IV) ports inserted "incanulation". Once inserted, healing medications, nutrients and fluids can be administered into the patient's bloodstream through the IV port. But the initial implantation of the IV port (e.g., sticking the IV port into the patient's arm) is often surprisingly painful. The same painful port insertion procedure is used with cancer patients, burn patients, patients being prepared for surgery, and a long list of other medical problems. And IV port placement is just one of large number of medical procedures where excessive pain is common.

Some medical procedures sound fairly painless, but patients' fears and anxiety associated with the procedures can amplify the patient's negative experience. For example, with blood draws, a fairly large percentage of people faint during blood draws, despite the short duration of the procedure.

A number of psychological influences can amplify or increase how much pain patients experience during medical procedures. High anxiety can increase sensitivity to pain. Some patients

DOI: 10.4018/978-1-5225-2255-3.ch518

must receive medical procedures repeatedly. For example, children with large severe burn injuries receive several painful medical procedures per week, often daily wound cleaning and painful physical therapy skin stretching exercises, and the most painful segments of the wound care procedures often last 25 minutes or longer. If the patient has an extremely high pain during wound care, their memory for previous painful wound care sessions can increase their subjective experience of pain during subsequent wound care procedures, and make it harder to control the patient's pain using pain medications alone. There are concerns that repeated painful medical procedures can be made worse if the patient has pre-existing psychological problems such as depression or PTSD. These conditions can reduce the effectiveness of opioid analgesics. And repeated painful medical procedures during hospitalization may increase patients' risk of developing chronic pain or chronic PTSD. There has lately been a growing realization in the medical community for the need for more effective non-pharmacologic pain control during medical procedures. Use of higher doses of pain medications may increase analgesia, but also increase undesirable medical side effects of the pain medications such as nausea, delirium, constipation, urinary retention, sleeping through meals, and interference with sleeping at night. Diversion of pain medications is another concern. On the streets of the U.S.A., addiction to opioids has become a major medical problem in itself. According to the CDC, overdose deaths involving prescription opioids have quadrupled since 1999, and set a new record high in 2014 (CDC, 2016).

Because pain has a strong psychological component, in theory, psychological interventions can be used to help counteract some of the negative psychological influences. In other words, there is growing interest in using psychological techniques to help reduce pain. Typically, psychological techniques can be used in addition to traditional pain medications, adjunctively (Stinson, Yamada, Dickson, Lamba & Stevens, 2008; Koller & Goldman, 2012). Distraction is a familiar technique. Parents frequently distract children to help get

them to stop crying (using a rattle toy). Music is widely used to help patients during dental procedures. According to a Cochrane meta-analysis, listening to music resulted in a statistically significant reduction in pain, but on average, music only reduced the amount of pain by around 5% reduction (not a clinically meaningful reduction from listening to music). A stronger distraction is needed for acute pain during medical procedures.

Recently Virtual Reality (VR) has emerged as a new unusually effective distraction technique, allowing the user to be immersed in a computer-generated environment. Patients look into virtual reality goggles that block the patient from being able to watch their wound care, and earphones and computer generated sound effects and music from VR hinder the sounds of the real environment, (Hoffman et al., 2000). The user can interact with the virtual environments using a mouse or a head-tracking.

Hoffman and Patterson and colleagues co-originated the technique of using immersive virtual reality analgesia during acute procedural pain (Hoffman, 1998; Hoffman et al, 2000). Two adolescents with severe burns used immersive virtual reality during burn wound care: staple removals. Both children reported large reductions in their pain during wound care while using immersive virtual reality, compared to when the same children played a Nintendo video game during the same wound care sessions (Hoffman et al., 2000). Several controlled studies support VR efficacy in reducing pain but most of them investigated its effects on adults, in particular during procedural pain in burn patients (Hoffman, Patterson, Magula, Carrougner, Zeltzer, Dagadakis & Sharar, 2004; Maani, Hoffman, Morrow, Maiers Gaylord, McGhee & DeSocio, 2011) and during dental procedures (Bentsen, Svensson & Wenzel, 2001; Hoffman, Garcia-Palacios, Patterson, Jensen, Furness III & Ammons, 2001; Furman, Jasinevicius, Bissada, Victoroff, Skillicorn & Buchner, 2009; Wiederhold, Gao & Wiederhold, 2014).

The aim of the chapter is to review the use of VR for pediatric and adolescent pain management and its applicability in clinical settings. Between

9 more pages are available in the full version of this document, which may be purchased using the "Add to Cart" button on the publisher's webpage:

www.igi-global.com/chapter/virtual-reality-as-distraction-technique-for-pain-management-in-children-and-adolescents/184296

Related Content

A Bio-Inspired, Distributed Control Approach to the Design of Autonomous Cooperative Behaviors in Multiple Mobile Robot Systems

Gen'ichi Yasuda (2018). *Encyclopedia of Information Science and Technology, Fourth Edition* (pp. 6836-6846).

www.irma-international.org/chapter/a-bio-inspired-distributed-control-approach-to-the-design-of-autonomous-cooperative-behaviors-in-multiple-mobile-robot-systems/184380

PRESCAN Adaptive Vehicle Image Real-Time Stitching Algorithm Based on Improved SIFT

Qian Li, Yanli Xuand Pengren Ding (2023). *International Journal of Information Technologies and Systems Approach* (pp. 1-17).

www.irma-international.org/article/prescan-adaptive-vehicle-image-real-time-stitching-algorithm-based-on-improved-sift/321754

Collaboration Network Analysis Based on Normalized Citation Count and Eigenvector Centrality

Anand Bihari, Sudhakar Tripathiand Akshay Deepak (2019). *International Journal of Rough Sets and Data Analysis* (pp. 61-72).

www.irma-international.org/article/collaboration-network-analysis-based-on-normalized-citation-count-and-eigenvector-centrality/219810

Efficient Techniques to Design Low-Complexity Digital Finite Impulse Response (FIR) Filters

David Ernesto Troncoso Romeroand Gordana Jovanovic Dolecek (2015). *Encyclopedia of Information Science and Technology, Third Edition* (pp. 1579-1589).

www.irma-international.org/chapter/efficient-techniques-to-design-low-complexity-digital-finite-impulse-response-fir-filters/112562

Pricing Based on Real-Time Analysis of Forklift Utilization Using RFID in Warehouse Management

Numan Celebi, Kübra Savaand Ihsan Hakan Selvi (2018). *Encyclopedia of Information Science and Technology, Fourth Edition* (pp. 5490-5502).

www.irma-international.org/chapter/pricing-based-on-real-time-analysis-of-forklift-utilization-using-rfid-in-warehouse-management/184251