Chapter 5 Demethylation of Arsenic and Nickel From Tannery Effluent

Shankar Mukundrao Khade Indian Institute of Technology (BHU), India

Suresh Kumar Srivastava Indian Institute of Technology (BHU), India

ABSTRACT

The tanning Industry is considered to be a major source of pollution and tannery wastewater in particular is a potential environmental concern. Wastewaters produced by tanneries contain high concentration of inorganic and organic pollutant. The pollutants of concern within the tanning industry include Azodyes, Cadmium compounds, Cobalt, Copper, Antimony, Barium, Lead, Selenium, Mercury, Zinc, Arsenic, Polychlorinated Biphyenls (PCB), Nickel, Formaldehyde resins, and Pesticides residues. Because tannery wastewater contains a complexity of pollutants including chromium and chlorinated phenols as indicated earlier, it is vital to dissect the toxic nature of such wastewater both to understand its environmental impacts and identify potential remediation strategies. Furthermore, there are strict regulations imposed for the environmental control of pollutants such as heavy metals and persistent organic pollutants. In this chapter, the authors discuss the demethylation process of Arsenic and Nickel from Tannery Effluent.

DOI: 10.4018/978-1-5225-4162-2.ch005

Copyright © 2018, IGI Global. Copying or distributing in print or electronic forms without written permission of IGI Global is prohibited.

INTRODUCTION

The setback of environmental pollution on report of vital industrial growth in practical requisites is the problem of disposal of industrial wastes in the form of solid, liquid, or gas. These forms are responsible for ultimately polluting water. The polluted water indeed affects the soil of industrial areas along with the fertile agricultural soils. In addition to these effects, secondary pollution is also prominent by mixing with the ground water. Industries are the one of the main cause of environmental pollution. The major of quantity and type of pollutants released into the environment depends on the type of industry. The effluent discharged from the industries cause severe metal pollution in the environmental. Most of the metals released into the environment cause severe risk to the human and animal's health. Many metals including arsenic, copper, mercury, chromium, cadmium, aluminium, nickel, zinc, and uranium affects the environment. These metals impose severe concerns to the organisms due to their persistence, biomagnification and bioaccumulation in food chain. So to avoid these effects, prior treatment of industrial waste is necessary. The exposure to these metals affects the health of an organism by retarding the development and growth, organ damage, neurological damages, cancer and at higher concentration may leads to death. The exposure to mercury leads to minamata syndrome. In addition to this, mercury and lead causes autoimmune disease along with kidneys, circulatory system (Barregård et al., 1999; Barsoum, 2006; Soderland et al., 2010).

The release of industrial effluents containing soluble or insoluble compounds into the ground water, streams, ponds, rivers, and oceans leads to cause primary and secondary water pollution (Akcil & Koldas, 2006; Gupta et al., 2012). However, the released pollutants are either removed or transformed to the less harmful compound by microbial approach termed as 'bioremediation'. Biotransformation is an enzymatic process which includes oxidation, reduction, methylation or demethylation of highly toxic pollutants. The microbes involved in maintaining the biogeochemical cycle of arsenic by demethylation process include *Candida humicola, Mycobacterium neoaurum, Bacillus sp.* etc. Most of the water pollution in the developing countries occurs because of the poor management of wastes generated in huge quantity due to many anthropogenic activities (Giusti, 2009; Goel, 2006).

ENVIRONMENTAL POLLUTION AND ITS EFFECTS

Globally environmental pollution is now days a big challenge, due to the usage of huge amount of chemicals and discharge of untreated industrial effluents into the water bodies. Heavy metals which are used in the various industries or produced 10 more pages are available in the full version of this document, which may be purchased using the "Add to Cart" button on the publisher's webpage: www.igi-

global.com/chapter/demethylation-of-arsenic-and-nickel-

from-tannery-effluent/204826

Related Content

Analysis of a Monopile under Two-Way Cyclic Loading

(2021). International Journal of Geotechnical Earthquake Engineering (pp. 0-0). www.irma-international.org/article//282687

Effective Waste Water Treatment and its Management

Sakthivel Lakshmana Prabu, TNK Suriyaprakash, Ruckmani Kandasamyand Thirumurugan Rathinasabapathy (2016). *Toxicity and Waste Management Using Bioremediation (pp. 312-334).*

www.irma-international.org/chapter/effective-waste-water-treatment-and-itsmanagement/141806

What Vietnamese Employers and Educators Want in Impactful Engineering Degree Programmes

Thuy Tien Nguyen, Anh Pham Thi, Tuan Anh Nguyen, Luu Mai, J. Toby Mottram, Georgia Kremmyda, Yiannis Koumpouros, Angelos Georgoulasand Dong Doan Van (2023). *Transcending Humanitarian Engineering Strategies for Sustainable Futures (pp. 1-20).*

www.irma-international.org/chapter/what-vietnamese-employers-and-educators-want-inimpactful-engineering-degree-programmes/320352

Spectral Estimation of Noisy Seismogram using Time-Frequency Analyses

Vaneeta Deviand M. L. Sharma (2016). *International Journal of Geotechnical Earthquake Engineering (pp. 19-32).*

www.irma-international.org/article/spectral-estimation-of-noisy-seismogram-using-time-frequency-analyses/174410

The Optimized Dynamic Behavior of Short Embankment Based on Frequency Performance

Behrouz Gordanand Azlan Adnan (2015). *International Journal of Geotechnical Earthquake Engineering (pp. 1-11).*

www.irma-international.org/article/the-optimized-dynamic-behavior-of-short-embankment-basedon-frequency-performance/159213