## Chapter 11 Newer Approaches in Phytoremediation: An Overview

## ABSTRACT

The heavy metal pollution problem is all over the world. Plant-growthpromoting bacteria (PGPB) has transformed heavy metals present in the soil, which removes and minimizes their toxic effects. This chapter highlights the role of plant-growth-promoting bacteria, chelating agents, and nanoparticles for remediation of heavy metals; their mechanism of action; and their applications approach of hyperaccumulation. Therefore, this chapter focuses on the mechanisms by which microorganisms, chelating agents, and nanoparticles can mobilize or immobilize metals in soils and the nano-phytoremediation strategies are addressed for the improvement of phytoextraction as an innovative process for enhancement of heavy metals removal from soil.

### INTRODUCTION

Heavy metals is a serious environmental problem which affecting human health and plant (Aldoobie and Beltagi, 2013; Park et al., 2014). The use of large amount of fertilizers has been produced heavy metals into non-pollluted sites which affect dramtically to agriculture (Saba et al., 2015). One of the most toxic heavy metal involved in many fertilizers is lead (Pb), which can be

DOI: 10.4018/978-1-5225-9016-3.ch011

absorbed and translocate into plants and easily enter into food chain. Heavy metal contaminants movement from soil to the ground water is very slow because of less mobility so it is not easily absorbed by the plant (Rodriguez et al., 2011; Lori et al., 2015). Therefore heavy metals are low in the upper parts such as leaves, seeds and fruits because of high amount of accumulation in roots (Mishra and dubey, 2005; Ullah et al., 2015). High concentration in soil can also affect to microbial activity with soil fertility (Gao et al., 2010; Yuan et al., 2015). Heavy metals such as Zn, Cu and Mn, Al, Mg are necessary as micronutrients, but high amount of these heavy metals can be caused toxicity and negative effect on human health (Langer et al., 2009; Ali et al., 2013). Some other heavy metals such as Cd, As, Pb, Hg and Ni, are non-nutritional and toxic elements present in soil (Jourand et al., 2010).

Plants under heavy metal stress condition it produces a high level of reactive oxygen species (ROS) such as hydrogen peroxide  $(H_2O_2)$ , hydroxyl radicals (OH), superoxide radicals (O<sub>2</sub>) catalse, which result in damages to plant cell or tissue (Wang et al., 2015; Migocka et al., 2014). Reactive oxygen species produced continuously in different compartments with antioxidant molecule under as by-product (Reddy et al., 2005; Kwankua et al., 2012; Girisha and Ragavendra, 2009; Palma et al., 2013; Dubey et al., 2014). However the critical imbalance and excess amount of production ROS, antioxidant molecule in plant which is depleted and creates disorders in plant enzymatic activity, all type of biochemical molecules including cell wall, membrane lipids, protein, amino acid chain and carbohydrates figure 1 (Kaur et al., 2015; Qiao et al., 2015; Gratao et al., 2005). They also damage cell membrane, loss of cell metabolites, reduction in cell growth and impaired metabolic functions (Goncalves et al., 2007a,b; Lee et al., 2007; Merlot et al., 2014; Morel et al., 2009).

The conventional remediation technologies are cost effective more expensive and some of the techniques did not remove heavy metal effectively. Therefore, it is important to develop economically practical and more effective method to decontaminate soils from heavy metal contamination. The most popular advantage of phytoremediation is low cost effectiveness (Abdul and Schroder, 2009). It can be up to 1000-fold cheaper compared to conventional method such as (flotation-filtration, evaporation, ion-exchange, electrodialysis and ultrafiltration). It has been estimated to clean up one acre of sandy loam soil to a depth of 55 cm will cost 60,000-100,000\$ compare to 400,000\$ for conventional using traditional soil removal methods (Ali et al., 2013).

32 more pages are available in the full version of this document, which may be purchased using the "Add to Cart" button on the publisher's webpage: <u>www.igi-</u> <u>global.com/chapter/newer-approaches-in-</u> <u>phytoremediation/241174</u>

### **Related Content**

#### Dairy Waste Treatment and Utilization: A Treatise

Anum Ishaq, Ubaid ur Rahman, Muhammad Haseeb Ahmad, Amna Sahar, Qamar Abbas Syedand Rashida Perveen (2020). *Waste Management Techniques for Improved Environmental and Public Health: Emerging Research and Opportunities (pp. 160-182).* 

www.irma-international.org/chapter/dairy-waste-treatment-and-utilization/243878

#### Physicochemical Characteristics of Indonesian Native Starch

Maria Belgis, Nur Fathonah Sadek, Ardiyan Dwi Masahid, Dian Purbasariand Dyah Ayu Savitri (2023). *Food Sustainability, Environmental Awareness, and Adaptation and Mitigation Strategies for Developing Countries (pp. 62-79).* www.irma-international.org/chapter/physicochemical-characteristics-of-indonesian-nativestarch/319451

### Remote Sensing-Based Evapotranspiration Modelling for Agricultural Water Management in the Limpopo Basin

Berhanu F. Alemaw, Thebeyame Ronald Chaokaand Brigton Munyai (2019). Environmental Information Systems: Concepts, Methodologies, Tools, and Applications (pp. 249-286).

www.irma-international.org/chapter/remote-sensing-based-evapotranspiration-modelling-foragricultural-water-management-in-the-limpopo-basin/212947

## Long Term Modeling and Predictive Assessment of Climate Change of Ukraine Polesie Zone

Petro Kovalenko, Nataliia Prykhodko, Vasyl Turcheniukand Jerzy Jeznach (2023). Handbook of Research on Improving the Natural and Ecological Conditions of the Polesie Zone (pp. 123-133).

www.irma-international.org/chapter/long-term-modeling-and-predictive-assessment-of-climatechange-of-ukraine-polesie-zone/324034

# Fabrication of Metal Oxide-Biopolymer Nanocomposite for Water Defluoridation

Dickson Mubera Andala, Erick Mobegi, Mildred Nawiriand Geoffrey Otieno (2020). Effects of Emerging Chemical Contaminants on Water Resources and Environmental Health (pp. 242-271).

www.irma-international.org/chapter/fabrication-of-metal-oxide-biopolymer-nanocomposite-forwater-defluoridation/248384