


Wildfire Air Quality Prediction: A Data-Driven Approach

Subhankar Dhar, San Jose State University, USA*

 <https://orcid.org/0000-0003-1493-451X>

Jerry Gao, San Jose State University, USA

ABSTRACT

Wildfires are extremely harmful to the environment. While producing gaseous pollutants and particles that cause smoke, wildfires also release carbon dioxide (CO₂), a greenhouse gas that will continue to warm the planet after the wildfire ends. This article delves into the impact of wildfires and air quality on human living conditions. The authors' machine learning models use wildfire data to forecast air quality with detailed indexes and geographic information during a wildfire. The work evaluates the performance of each machine learning model via statistical metrics like mean absolute error (MAE), R-squared (R²), and root mean squared error (RMSE). The experimental results used neural networks to predict a specific value for carbon monoxide (CO), ozone, and PM_{2.5}. These are both promising and accurate, providing meaningful insight into air quality within a region. This work will be useful for cities, governments, citizens, and public safety.

KEYWORDS

Air Pollution, Air Quality Prediction, Big Data Analytics, Disaster Response Management, Environment Management, Machine Learning, Neural Network, Wildfires

INTRODUCTION

Wildfires can increase air pollution and cause severe damage to air quality by emitting carbon dioxide (CO₂), carbon monoxide (CO), and other greenhouse gases that contribute to global warming and environmental hazards (Castelli et al., 2020). In addition, they can damage forests that otherwise remove CO₂ from the air and inject aerosols into the atmosphere. Smoke from wildfires can cause serious health disorders and respiratory diseases (Camia & Amatulli, 2009; de Groot et al., 2007) by reducing the size of green forests and eliminating CO₂ in the air (Reid et al., 2019). Pollutants like particulate matter (PM) 2.5, ozone (O₃), nitrogen dioxide (NO₂), and CO are major parameters in the air quality index (AQI) of a region (Tian et al., 2011).

DOI: 10.4018/IJDREM.330148

*Corresponding Author

This article published as an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0/>) which permits unrestricted use, distribution, and production in any medium, provided the author of the original work and original publication source are properly credited.

This work delves into the impact of wildfires on air quality, using data science tools and technologies to predict the impact of air quality after a wildfire event. To achieve this goal, the authors must define key measurements with respect to air. For example, air has several important measurement metrics that affect air quality, including the CO level, ozone level, and PM2.5 (small inhalable particles with diameters no more than 2.5 micrometers) index. These pollutants can have severe negative impacts on human health, including wheezing, coughing, sore eyes, and throats. The authors, therefore, develop machine learning models to understand the health effects caused by wildfires and strengthen public awareness surrounding such events.

First, the authors develop a data-driven machine learning framework to predict air quality during and after wildfires. In addition, the study addresses the limitations of existing methods. The research provides insight into the significant impact of wildfires, enabling people and governments to prioritize environmental protections and prevention efforts. Residents can also monitor air quality changes more closely and prepare for potential wildfires.

Second, the authors use two sets of machine learning models to predict air quality during wildfire seasons. The study provides detailed explanations on how to determine and tune parameters. Analysis helps the public understand and anticipate air quality changes. It also enables individuals to compare and improve upon existing models.

Third, the study includes results driven by statistical analysis and machine learning. It aims to prove the significance of wildfires and predict air quality with improved accuracy. Descriptive and inferential statistical analysis tools, along with visualization diagrams and graphs, depict the difference between air quality before and after wildfires. The study's regression and deep learning models generate precise metrics to increase accuracy.

Fourth, the study presents a data-driven approach for deploying an online wildfire air quality alert system. It uses real-life data to develop a proof-of-concept implementation. This prototype demonstrates the novel functionalities of the authors' framework, benefiting both the public and city officials.

The article is organized as follows. The next section discusses related works. Then, the article describes the geographical location of the study, comprehensive data parameters, data pre-processing, and training and test data preparation. Wildfire air quality prediction models and experimental results are explored before navigating the system design, implementation framework, and visualization tools. Lastly, the study discusses the conclusion and future work.

RELATED WORK

Preisler et al. (2015) highlighted the shortcomings of research when predicting wildfire impacts from PM2.5 concentration at ground-level monitors in California. While most researchers rely on satellite-based observational tools, this work combined models with an autoregressive statistical model, incorporating weather and seasonal factors to identify thresholds for predicting unusual events. The study focused on ground-based monitoring of PM2.5 levels, with data consisting of hourly values of PM2.5 and meteorological data. Data was gathered from the United States Department of Agriculture Forest Services. Unexpectedly, the study found that smoke plumes could identify seasonal wildfire influence with high accuracy.

Reid et al. (2015) and Jaffe et al. (2013) evaluated the contribution of O₃ caused by wildfires in the atmosphere in western U.S. The studies developed several statistical models to estimate the maximum daily eight-hour average (MDA8) O₃ emissions and created methodology functions for three western areas in the U.S. The residual of the statistical model can provide information on O₃ emissions that cannot be explained by normal wildfires.

Similarly, Preisler and Westerling (2007) developed a statistical model for forecasting fire-danger and producing one-month ahead wildfire-danger probability in the western U.S. The model predicted a monthly average temperature and drought severity index to demonstrate significant potential

20 more pages are available in the full version of this document, which may be purchased using the "Add to Cart" button on the publisher's webpage: www.igi-global.com/article/wildfire-air-quality-prediction/330148

Related Content

Integrating Volunteers into Rescue Processes: Analysis of User Requirements and Mobile App Conception

Henrik Detjen, Stefan Hoffmann, Leonie Rösner, Stephan Winter, Stefan Geisler, Nicole Krämer and Gerd Bumiller (2015). *International Journal of Information Systems for Crisis Response and Management* (pp. 1-18).

www.irma-international.org/article/integrating-volunteers-into-rescue-processes/143918

Communicating Location and Geography in Emergency Response

Fredrik Bergstrand, Jonas Landgren and Urban Nuldén (2019). *Emergency and Disaster Management: Concepts, Methodologies, Tools, and Applications* (pp. 1048-1066).

www.irma-international.org/chapter/communicating-location-and-geography-in-emergency-response/207615

Economics and Practical Applications for Applied Trauma Theory: Sustainable Energy and Rural Tourism

David S. Bathory (2014). *Crisis Management: Concepts, Methodologies, Tools, and Applications* (pp. 1222-1236).

www.irma-international.org/chapter/economics-and-practical-applications-for-applied-trauma-theory/90774

Review of Agent Based Modelling of Social Attachment in Crisis Situations

Julius M. Bañgate, Julie Dugdale, Elise Beck and Carole Adam (2019). *International Journal of Information Systems for Crisis Response and Management* (pp. 35-64).

www.irma-international.org/article/review-of-agent-based-modelling-of-social-attachment-in-crisis-situations/234326

A Proposed Framework for Developing a National Crisis Management Information System

Magdy M. Kabeil (2009). *International Journal of Information Systems for Crisis Response and Management* (pp. 50-74).

www.irma-international.org/article/proposed-framework-developing-national-crisis/37526