


Automated Detection of On-Farm Irrigation Reservoirs in Two Critical Groundwater Regions of Arkansas: A Necessary Precursor for Conjunctive Water Management


Daniel D. Shults, Arkansas State University, USA

John W. Nowlin, Arkansas State University, USA*

 <https://orcid.org/0000-0002-2891-028X>

Joseph H. Massey, USDA-ARS, USA

Michele L. Reba, USDA-ARS, USA

 <https://orcid.org/0000-0001-6830-0438>

ABSTRACT

In eastern Arkansas, the use of surface water for crop irrigation is steadily increasing in response to declining aquifers. Effective conjunctive water management requires accurate and timely information on the locations, sizes, and numbers of on-farm irrigation reservoirs. A method for remotely locating and characterizing on-farm reservoirs was developed using relative elevation and near-infrared imagery. With 62% accuracy, the method automatically identified 429 irrigation reservoirs within a 1.9-Mha area in less than an hour using an off-the-shelf laptop. Reservoirs not accurately identified (i.e., false negatives) were caused by the presence of vegetation or turbidity within the reservoirs. There were no false positive detections. This approach for identifying elevated reservoirs is applicable across the Mississippi Alluvial Plain (MAP) that encompasses over 4-Mha of irrigated cropland and other agricultural areas having low-relief.

KEYWORDS

ArcGIS Pro Model Builder, Automated Reservoir Detection, Conjunctive Water Management, Topographic Modeling, Waterbody Classification

INTRODUCTION

The Mississippi River Valley Alluvial Aquifer (MRVAA) provides 90% of the irrigation water applied to crops in the Mississippi Alluvial Plain (MAP) (Leslie et al., 2022). The aquifer has been over-exploited for decades (Bedinger et al., 1964; Clark et al., 2011; 2013; Vories & Evett, 2014). In order to capture precipitation/runoff and reduce groundwater withdrawals, irrigation reservoirs are being

DOI: 10.4018/IJAGR.337287

*Corresponding Author

This article published as an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0/>) which permits unrestricted use, distribution, and production in any medium, provided the author of the original work and original publication source are properly credited.

constructed on farms throughout the MAP (Evelt et al., 2003). Additionally, in central Arkansas, two large surface water diversion projects will use on-farm reservoirs to store water removed from the White River and the Arkansas River to offset groundwater pumping (USACE, 1999; 2007).


These on-farm reservoirs can be stand-alone structures or may be part of more complex systems having a tailwater recovery ‘pit’ or ‘ditch’ to capture runoff (Figure 1). In both cases, the reservoirs are generally symmetrical (e.g., square, rectangular, triangular), relatively compact (i.e., small perimeter-to-surface area ratio), and surrounded by raised earthen levees (Yaeger et al., 2018).

Strategies to conjunctively manage surface and ground water resources (Kovacs et al., 2016; Singh et al., 2016) have been adopted in Arkansas (ANRC, 2014; 2016; 2017; 2019), Mississippi (YMD, 2006), Louisiana (E&E INC, 2011), and Missouri (MDNR, 2020). As a result, accurate and up-to-date information on irrigation reservoirs is necessary to allow resource managers and policymakers to (a) assess how existing reservoirs may impact groundwater demand, (b) determine where surface waters could support additional reservoir(s), (c) estimate maintenance costs and life expectancies of existing reservoir infrastructure, and (d) determine carrying capacity of reservoir area per watershed area.

Figure 1 shows examples of (A) two on-farm irrigation reservoirs in Eastern Arkansas that receive water pumped from streams and/or a tailwater recovery ditch (center) and (B) corresponding elevation profile. Note the western reservoir’s high turbidity in contrast to the eastern reservoir.

The construction of reservoirs in the MAP may occur solely at the expense of farmers and/or landowners but more commonly occurs with assistance from federal and state agencies (Czarnecki et al., 2024; King, 2021). In Arkansas, approximately \$45 million were awarded by the USDA Natural

Figure 1. Irrigation reservoirs in eastern arkansas and their corresponding elevation profiles



20 more pages are available in the full version of this document, which may be purchased using the "Add to Cart" button on the publisher's webpage: www.igi-global.com/article/automated-detection-of-on-farm-irrigation-reservoirs-in-two-critical-groundwater-regions-of-arkansas/337287

Related Content

GML-Based Data Management and Semantic World Modelling for a 4D Forest Simulation and Information System

Jürgen Roßmann, Martin Hoppenand Arno Bücken (2014). *International Journal of 3-D Information Modeling* (pp. 50-67).

www.irma-international.org/article/gml-based-data-management-and-semantic-world-modelling-for-a-4d-forest-simulation-and-information-system/122867

Studying Surface and Canopy Layer Urban Heat Island at Micro-Scale Using Multi-Sensor Data in Geographic Information Systems

Bakul Budhiraja, Prasad Avinash Pathakand Debopam Acharya (2019). *Geospatial Intelligence: Concepts, Methodologies, Tools, and Applications* (pp. 389-410).

www.irma-international.org/chapter/studying-surface-and-canopy-layer-urban-heat-island-at-micro-scale-using-multi-sensor-data-in-geographic-information-systems/222908

Reinforcing the Place of Dynamic Spatialised Indicators in a Generic Socioeconomic Model

Fenintsoa Andriamasinoro (2013). *Geographic Information Analysis for Sustainable Development and Economic Planning: New Technologies* (pp. 313-334).

www.irma-international.org/chapter/reinforcing-place-dynamic-spatialised-indicators/69065

Local Perspectives on Spatial Accessibility to Market in the Afram Plains, Ghana

Nathaniel Dede-Bamfo, T. Edwin Chowand Chikodinaka Vanessa Ekeanyanwu (2023). *International Journal of Applied Geospatial Research* (pp. 1-24).

www.irma-international.org/article/local-perspectives-on-spatial-accessibility-to-market-in-the-afram-plains-ghana/322409

Methodical Spatial Database Design with Topological Polygon Structures

Jean Damascène Mazimpaka (2012). *International Journal of Applied Geospatial Research* (pp. 21-30).

www.irma-international.org/article/methodical-spatial-database-design-topological/62044