

# Modeling: A Central Activity for Flexible Information Systems Development in Agriculture and Environment

*P. Papajorgji, University of Florida, USA*

*F. Pinet, Cemagref, Clermont Ferrand, France*

*A. Miralles, Cemagref, Maison de la Télédétection, France*

*E. Jallas, ITK/CIRAD, France*

*P.M. Pardalos, University of Florida, USA*

---

## ABSTRACT

*Enterprise information systems in agriculture and environment are becoming more complex and difficult to design and implement. This paper aims to show our vision on using model-based approaches to design complex and flexible agricultural and environmental information systems. At the center of this modeling approach is the Unified Modeling Language that facilitates expressing visually concepts of a problem domain and their relationships. UML has a core of notations that are generic and that can be used to model problems in any domain but can be extended to create profiles in order to take into consideration modeling concerns in a particular problem domain. UML profiles are created to use UML in designing spatial systems, ontologies, model driven architecture-based systems and Web-based systems and a recent profile makes it possible to use UML for business modeling purposes. UML is used to present design patterns; their use is crucial in designing complex and flexible information systems. Recently, UML is enriched with Object Constraint Language that is used to express constraints on modeling artifacts. The paper presents the state of the art in modeling agricultural and environmental systems and provides discussions for future directions.*

**Keywords:** *Business Applications, Environmental Modeling, Object Oriented Design, Ontologies, Spatial Data, Unified Modeling Language*

---

## INTRODUCTION

During the past decades agriculture has experienced substantial structural and technological

changes. Currently, agricultural production is organized around large farms that can afford applying new technologies. At the same time, social consciousness about environmental issues is very strong. Land management is a complex process and to manage it successfully it requires that special attention should be paid

DOI: 10.4018/jaeis.2010101301

Figure 1. Software development process (Jacobson, Booch, & Rumbaugh, 1999).



to environmental issues while designing and developing agricultural and environmental information systems.

On the one hand, the advent of the Internet has opened vast opportunities for communication and business opportunities within the agricultural and environmental communities. On the other hand, it has substantially increased the amount of information farmers and territory managers need to analyze and synthesize to successfully administrate all the multi-facets of the agricultural production and land management. This new complex environment requires new and complex approaches to carry on farm production and land governance. Complex systems are not easy to design and develop unless a rigorous methodology and the right tools are used. Therefore, it is imperative that modern software engineering approaches be used in agriculture and environment-related areas.

Since the earliest days of software engineering, many methodologies have been designed and used to create better software. An important event that significantly influenced the software engineering world was the use of visual modeling tools. Modeling is a well-known engineering discipline as it helps one to understand reality. Models of complex systems are built because it is difficult to understand such a system in its entirety. Models are needed to express the structure and the behavior of complex systems. Using models makes it possible to visualize and better control system's architecture (Papajorgji & Shatar, 2004).

In the early 90s there were several modeling languages used by the software engineering community. The most well-known methodologies were Booch, Jacobson, (Object-Oriented Software Engineering) and Rumbaugh's (Object Modeling Technique). Other important methods

were Fusion (Coleman et al., 1994), Shlaer-Mellor (Shlaer & Mellor, 1988), and Coad-Yourdon (Coad & Yourdon, 1991). All these methods had different strengths and weaknesses. An important event occurred in the mid 1990s when Booch, Jacobson, and Rumbaugh began adopting ideas from each other that led to the creation of the Unified Modeling Language or as it is known best, the UML (OMG, 2009b). Details about UML will be presented.

They define a software development process as the set of activities needed to transform a user's requirements into a software system (see Figure 1).

The various "ingredients" needed in the process of developing computer applications are presented in Figure 2. This model is called the 4 Ps model (Jacobson, Booch, & Rumbaugh, 1999). This model shows that the *Result* of a *Project* is a *Product* which requires different types of ingredients:

- *People* (actors) in order to describe the analyzed domain and *People* to develop and manage the *Project* (analysts, designers, programmers, etc.).
- A method (*Process*) to conduct the development following a framework (*Templates*) defining, organizing and explaining the successive steps of the development.
- *Tools* to facilitate the expression of the needs, the modeling, the project planning, the programming, and so forth.

Another important event that shaped the way software engineering was carried out is the design patterns movement. Design patterns are well-thought solutions for a large number of

23 more pages are available in the full version of this document, which may be purchased using the "Add to Cart" button on the publisher's webpage: [www.igi-global.com/article/modeling-central-activity-flexible-information/39025](http://www.igi-global.com/article/modeling-central-activity-flexible-information/39025)

## Related Content

---

### Demethylation of Arsenic and Nickel From Tannery Effluent

Shankar Mukundrao Khade and Suresh Kumar Srivastava (2022). *Research Anthology on Emerging Techniques in Environmental Remediation* (pp. 508-516). [www.irma-international.org/chapter/demethylation-of-arsenic-and-nickel-from-tannery-effluent/291253](http://www.irma-international.org/chapter/demethylation-of-arsenic-and-nickel-from-tannery-effluent/291253)

### Nitrate, Total Ammonia, and Total Suspended Sediments Modeling for the Mobile River Watershed

Vladimir J. Alarcon and Gretchen F. Sassenrath (2017). *International Journal of Agricultural and Environmental Information Systems* (pp. 20-31). [www.irma-international.org/article/nitrate-total-ammonia-and-total-suspended-sediments-modeling-for-the-mobile-river-watershed/179581](http://www.irma-international.org/article/nitrate-total-ammonia-and-total-suspended-sediments-modeling-for-the-mobile-river-watershed/179581)

### On the Use of Abduction as an Alternative to Decision Trees in Environmental Decision Support Systems

Franz Wotawa (2011). *International Journal of Agricultural and Environmental Information Systems* (pp. 63-82). [www.irma-international.org/article/use-abduction-alternative-decision-trees/51632](http://www.irma-international.org/article/use-abduction-alternative-decision-trees/51632)

### Biodegradation of Low Density Polyethylene Films

Soumita Dutta Laha, Kingshuk Dutta and Patit Paban Kundu (2018). *Handbook of Research on Microbial Tools for Environmental Waste Management* (pp. 282-318). [www.irma-international.org/chapter/biodegradation-of-low-density-polyethylene-films/206536](http://www.irma-international.org/chapter/biodegradation-of-low-density-polyethylene-films/206536)

### Modeling and Operating Strategies of Micro-Grids for Renewable Energy Communities

Saad Salman Khan (2016). *Smart Grid as a Solution for Renewable and Efficient Energy* (pp. 97-122). [www.irma-international.org/chapter/modeling-and-operating-strategies-of-micro-grids-for-renewable-energy-communities/150317](http://www.irma-international.org/chapter/modeling-and-operating-strategies-of-micro-grids-for-renewable-energy-communities/150317)