

Chapter 3.11

Semi- E -Preinvex Functions

Yu-Ru Syau

National Formosa University, Taiwan

E. Stanley Lee

Kansas State University, USA

ABSTRACT

A class of functions called semi- E -preinvex functions is defined as a generalization of semi- E -convex functions. Similarly, the concept of semi- E -quasiconvex functions is also generalized to semi- E -prequasiinvex functions. Properties of these proposed classes are studied, and sufficient conditions for a nonempty subset of the n -dimensional Euclidean space to be an E -convex or E -invex set are given. The relationship between semi- E -preinvex and E -preinvex functions are discussed along with results for the corresponding nonlinear programming problems.

1. INTRODUCTION

Convexity plays an important role in optimization theory. Various generalizations of convex functions have appeared in the literature. A significant generalization of convex functions is the introduction of preinvex functions, originated by (Hanson and Mond, 1987) but so named by (Jeyakumar, 1985). Later, (Weir and Mond, 1988) studied how and where preinvex functions can replace convex functions in multiple objective optimization.

Recently, (Youness, 1999) introduced the concepts of E -convex sets and E -convex functions

which generalize the definitions of convex sets and convex functions based on the effecting of an operator E on the domain on which functions are defined. Youness's initial results inspired a great deal of subsequent work, see (Duca et al., 2000; Duca & Lupsa, 2006; Fulga & Preda, 2009; Syau & Lee, 2005; Yang, 2001). In an earlier paper (Syau & Lee, 2005), we extended the class of E -convex functions to E -quasiconvex functions. More recently, (Fulga and Preda, 2009) introduced a class of sets, called E -invex sets, which generalize the concept of E -convex sets. Based on the proposed E -invex sets, they introduced the class of E -preinvex functions as a generalization of E -convex and preinvex functions, and extended the

DOI: 10.4018/978-1-61350-456-7.ch3.11

class of E -quasiconvex functions to E -prequasiinvex functions.

Recall that the lower level sets of convex functions are convex sets, and that the lower level sets of preinvex functions are invex sets. Due to these properties, a convex (resp. preinvex) programming with inequality constraints may be easily formulated. However, an E -convex function does not guarantee E -convexity of its lower level sets. Based on the concept of E -convex sets, (Chen, 2002) introduced and studied a new concept of semi- E -convex functions which guarantee E -convexity of their lower level sets.

Motivated both by earlier research works (Chen, 2002; Fulga & Preda, 2009) and by the importance of preinvexity, we introduce a class of functions called semi- E -preinvex functions which are generalizations of semi- E -convex functions. Similarly, the concept of semi- E -quasiconvex functions is also generalized to semi- E -prequasiinvex functions. Some properties of these proposed classes are studied, and sufficient conditions for a nonempty subset of the n -dimensional Euclidean space to be an E -convex or E -invex set are given. The relationship between semi- E -preinvex and E -preinvex functions is discussed. In addition, some results for the corresponding nonlinear programming problems are considered.

2. PRELIMINARIES

Let R^n denote the n -dimensional Euclidean space. Recall (Youness, 1999) that, by definition, a set $M \subseteq R^n$ is said to be E -convex if there is a mapping $E: R^n \rightarrow R^n$ such that

$$\lambda E(x) + (1 - \lambda) E(y) \in M, \forall x, y \in M, \forall \lambda \in [0, 1].$$

Let $E: R^n \rightarrow R^n$ be a given mapping. For a nonempty set $S \subseteq R^n$, let

$$E(S) = \{E(x): x \in S\}.$$

Lemma 2.1 (Youness, 1999, Proposition 2.2).

Let $M \subseteq R^n$ be a nonempty E -convex set, then $E(M) \subseteq M$.

A set $K \subseteq R^n$ is said to be an invex set with respect to (w.r.t. in short) a given mapping $\eta: R^n \times R^n \rightarrow R^n$ if

$$x, y \in K, \lambda \in [0, 1] \Rightarrow y + \lambda \eta(x, y) \in K.$$

Definition 2.1 (Mohan & Neogy, 1995). Let $K \subseteq R^n$ be a nonempty invex set w.r.t. a given mapping $\eta: R^n \times R^n \rightarrow R^n$. A function $f: K \rightarrow R^1$ is said to be preinvex on K w.r.t. η if for all $x, y \in K$ and $\lambda \in [0, 1]$,

$$f(y + \lambda \eta(x, y)) \leq \lambda f(x) + (1 - \lambda) f(y).$$

Throughout this section, let $\eta: R^n \times R^n \rightarrow R^n$ be a fixed mapping.

Definition 2.2 (Chen, 2002; Syau & Lee, 2005; Youness, 1999). Let $M \subseteq R^n$ be nonempty. A function $f: R^n \rightarrow R^1$ is said to be

E -convex on M if there is a mapping $E: R^n \rightarrow R^n$ such that M is an E -convex set and

$$f(\lambda E(x) + (1 - \lambda) E(y)) \leq \lambda f(E(x)) + (1 - \lambda) f(E(y))$$

for all $x, y \in M$ and $\lambda \in [0, 1]$;

E -quasiconvex on M if there is a mapping $E: R^n \rightarrow R^n$ such that M is an E -convex set and

$$f(\lambda E(x) + (1 - \lambda) E(y)) \leq \max \{f(E(x)), f(E(y))\}$$

for all $x, y \in M$ and $\lambda \in [0, 1]$; semi- E -convex on M if there is a mapping $E: R^n \rightarrow R^n$ such that M is an E -convex set and

$$f(\lambda E(x) + (1 - \lambda) E(y)) \leq \lambda f(x) + (1 - \lambda) f(y)$$

for all $x, y \in M$ and $\lambda \in [0, 1]$; semi- E -quasiconvex on M if there is a mapping $E: R^n \rightarrow R^n$ such that M is an E -convex set and

5 more pages are available in the full version of this document, which may be purchased using the "Add to Cart" button on the publisher's webpage:

www.igi-global.com/chapter/semi-preinvex-functions/62471

Related Content

Online Testing of Nondeterministic Systems with the Reactive Planning Tester

Jüri Vain, Marko Kääramees and Maili Markvardt (2012). *Dependability and Computer Engineering: Concepts for Software-Intensive Systems* (pp. 113-150).

www.irma-international.org/chapter/online-testing-nondeterministic-systems-reactive/55327

Bridging the 15 Million Person Mentoring Gap

Caroline Kim Ohand Theresa Stroisch (2012). *Computer Engineering: Concepts, Methodologies, Tools and Applications* (pp. 1723-1732).

www.irma-international.org/chapter/bridging-million-person-mentoring-gap/62540

Applying Formal Methods to Verify Web Services Orchestration and Choreography

Zohra Sbati (2018). *Multidisciplinary Approaches to Service-Oriented Engineering* (pp. 221-243).

www.irma-international.org/chapter/applying-formal-methods-to-verify-web-services-orchestration-and-choreography/205301

Engaging Software Engineering Students With Employability Skills

Jocelyn Armarego (2018). *Computer Systems and Software Engineering: Concepts, Methodologies, Tools, and Applications* (pp. 1800-1838).

www.irma-international.org/chapter/engaging-software-engineering-students-with-employability-skills/192948

Affective Facial Expressions Using Auto-Associative Neural Network in Kansei Robot “Ifbot”

Masayoshi Kanoh, Tsuyoshi Nakamura, Shohei Kato and Hidenori Itoh (2011). *Kansei Engineering and Soft Computing: Theory and Practice* (pp. 215-236).

www.irma-international.org/chapter/affective-facial-expressions-using-auto/46400