Chapter 21 High-Performance Computing for Theoretical Study of Nanoscale and Molecular Interconnects

Rasit O. Topaloglu GlobalFoundries, USA

Swati R. Manjari Rensselaer Polytechnic Institute, USA

Saroj K. Nayak Rensselaer Polytechnic Institute, USA

ABSTRACT

Interconnects in semiconductor integrated circuits have shrunk to nanoscale sizes. This size reduction requires accurate analysis of the quantum effects. Furthermore, improved low-resistance interconnects need to be discovered that can integrate with biological and nanoelectronic systems. Accurate system-scale simulation of these quantum effects is possible with high-performance computing (HPC), while high cost and poor feasibility of experiments also suggest the application of simulation and HPC. This chapter introduces computational nanoelectronics, presenting real-world applications for the simulation and analysis of nanoscale and molecular interconnects, which may provide the connection between molecules and silicon-based devices. We survey computational nanoelectronics of interconnects and analyze four real-world case studies: 1) using graphical processing units (GPUs) for nanoelectronic simulations; 2) HPC simulations of current flow in nanotubes; 3) resistance analysis of molecular interconnects and subject in transport improvement in graphene interconnects. In conclusion, HPC simulations are promising vehicles to advance interconnects and study their interactions with molecular/ biological structures in support of traditional experimentation.

DOI: 10.4018/978-1-4666-5125-8.ch021

INTRODUCTION

Silicon technology and nanoelectronics have provided vast opportunities to advance the technology, including the advances in supercomputers and stream processors¹. While the International Technology Roadmap for Semiconductors (ITRS, 2010) predictions suggest a silicon future of at least ten more years, the device, interconnect, or process technology to be used to advance us towards such predictions is not rigidly defined. Devices with gate lengths as small as few nanometers are actively researched, though it is not certain whether much smaller devices will be connected with each other with copper interconnects, or will indeed require integration with nanotubebased interconnects. Furthermore, requirements for low-cost and small medical devices, and biointegrated sensors stimulate researchers towards integrating nanoelectronics with molecules (Birge, 1994; Stan et al., 2003). It has become indispensable to utilize supercomputers, which were manufactured through the utilization of the latest silicon-technology, to further the lifespan of silicon and integrate or replace copper interconnects with *nanotubes*, nanowires, or *molecular* interconnects towards the realization of systems that are low cost and error resilient.

Electronic characterization of devices of such small scale requires analysis and understanding of many nanoscale effects. These nanoscale effects, such as surface scattering² and quantum confinement³, are related to atomic interactions. For a given effect, the number of atoms involved in a region of a device can exceed thousands. Due to the nanoscale nature of the problem, effects that were not considered appreciable, such as quantum capacitance and contact resistance (Wang et al., 2009), may actually become important. While techniques such as continuum simulations, where groups of atoms are treated as one continuous medium, are possible for certain problems, most accurate results can be obtained through trulyatomic simulations.

At nanometer scales, electronic structure-based simulations are frequently used. There are two aspects to such simulations. First, the physics behind the molecule placement and interactions needs to be solved. This involves solving the Schrödinger equation, for which an exact analytic solution is not available for most systems. The solution can be quite complex. The next aspect is the visualization of molecular structures or the obtained solutions. Real-time, high-speed, and realistic visualization enables analysis and understanding of the physical effects. Both of these aspects require high-performance computing (HPC) for feasible turnaround time.

There are many HPC tools in the area of molecular dynamics. The complexity and implemented methods have been improved over time. Some examples of tools used for classical MD simulations are CHARMM (Brooks et al., 1982), LAMMPS (Sandia National Laboratories, 2011; Plimpton, 1995), NAMD (Phillips et al., 2005), and GROMACS (Van der Spoel et al., 2005; Hess et al., 2008) to name a few. Another molecular dynamics software package Desmond (Desmond, 2011), developed recently by D. E. Shaw Research, provides high-speed molecular dynamics simulations for biological systems that are highly scalable with the number of processors on conventional compute clusters (Bowers et al., 2006). However, when Desmond is used on the special-purpose supercomputer Anton, also designed by D. E. Shaw Research, speeds that are orders of magnitude faster can be achieved (Shaw et al., 2009). For the visualization aspect, Visual Molecular Dynamics package (VMD) (University of Illinois at Urbana-Champaign, 2011) is widely used. Advances are continually being made to improve existing visualization tools by using novel HPC methodologies and algorithms (Stone et al., 2009). In this chapter, we provide examples of more recent molecular dynamics and electronic transport simulation packages. Next, we give an overview of semiconductor interconnects and how high-performance nanoelectronic computa18 more pages are available in the full version of this document, which may be purchased using the "Add to Cart" button on the publisher's webpage:

www.igi-global.com/chapter/high-performance-computing-for-theoreticalstudy-of-nanoscale-and-molecular-interconnects/102027

Related Content

Synthesis and Characterization of Carbon-Based Quantum Dot From Rice, Sugar, and Aloe-Vera

Sunil Kumar Pradhan, Akula Maheswari Durga, Sumit Kashyapand Jostin Vara (2023). *Diversity and Applications of New Age Nanoparticles (pp. 321-335).*

www.irma-international.org/chapter/synthesis-and-characterization-of-carbon-based-quantum-dot-from-rice-sugar-andaloe-vera/321051

Bacterial Exopolysaccharides-Based Nanomaterials for Targeting Biofilm-Associated Infections: Bacterial Nanomaterials

Anand Ravichandran, D. Shobana, S. Visudha, Yazhini, S. Nivedithaand P. Saravanan (2024). *Cutting-Edge Applications of Nanomaterials in Biomedical Sciences (pp. 372-386).*

www.irma-international.org/chapter/bacterial-exopolysaccharides-based-nanomaterials-for-targeting-biofilm-associatedinfections/336405

Preparation and Application of Polymer-Metal Oxide Nanocomposites in Wastewater Treatment: Challenges and Potentialities

Victor Odhiambo Shikukuand Wilfrida N. Nyairo (2021). Research Anthology on Synthesis, Characterization, and Applications of Nanomaterials (pp. 393-407).

www.irma-international.org/chapter/preparation-and-application-of-polymer-metal-oxide-nanocomposites-in-wastewatertreatment/279158

DNA-Based Indexing

Max H. Garzon, Kiran C. Bobba, Andrew Neeland Vinhthuy Phan (2010). *International Journal of Nanotechnology and Molecular Computation (pp. 25-45).* www.irma-international.org/article/dna-based-indexing/52087

Nanotechnology Innovation Systems: A Regional Comparison

Nazrul Islam (2010). International Journal of Nanotechnology and Molecular Computation (pp. 65-84). www.irma-international.org/article/nanotechnology-innovation-systems/48529