

# Chapter 11

## New Imaging and Computational Technology as a Guide for Catheter Ablation of Incessant Tachyarrhythmias

**Carlo Pappone**  
University of Milan, Italy

**Vincenzo Santinelli**  
University of Milan, Italy

**Carmine Garzillo**  
University of Naples, Federico II, Italy

**Simonetta Crisà**  
University of Milan, Italy

### ABSTRACT

*Computational technology in the era of catheter ablation (RFA) has made it possible to experience relief from incessant atrial tachyarrhythmias (AT) by 3D electroanatomical mapping (EAM) systems. The Authors report the results of such technology in > 500 consecutive patients (57% males, mean age 56.9 years) with incessant refractory post-ablation left AT (mean cycle length 256 ms). Patients underwent electroanatomical-mapping systems, which combine electrophysiological and spatial information allowing accurate reconstruction of the whole atria with real-time activation sequence guiding RFA for continuous transmural linear lesions. Color-coded voltage and/or activation maps were successfully performed in all patients. Mapping distinguished clearly and rapidly between micro-macro-reentrant (>80%) and focal mechanisms. Acute success was obtained without major complications, with repeated procedures in about 5% of patients. EAM technology allows determining both mechanism and location of arrhythmia, ensuring successful elimination of complex arrhythmogenic substrates.*

### INTRODUCTION

#### An Epochal Change in the Management of Complex Tachyarrhythmias: Background

In the last two decades the rapid expansion of indications for catheter ablation from common supraventricular tachycardias to very complex tachyarrhythmias including atrial fibrillation, incessant atrial tachycardia and ventricular tachycardia led electrophysiologists to face more prolonged procedure times

DOI: 10.4018/978-1-5225-0140-4.ch011

with excessive fluoroscopy exposure and, then the need for stable and reproducible catheter movement, all of which require a substantial improvement in the existing traditional 2D mapping technology, and new developments (Silka, Gillette, et al., 1985; Van Hare, Velvis, & Langberg, 1990; Kay, Chong, et al., 1993; Chen, Chiang, et al., 1994; Lesh, Van Hare, et al., 1994; Poty, 1996; Chen, Tai, et al., 1998; Haissaguerre, Jais, et al., 1998; Schilling, Peters, Davies, 1998; Gornick, Adler, et al., 1999; Jais, Shah, et al., 2000; Kottkamp, Hugl, et al., 2000; Willems, Weiss, et al., 2000; Saoudi, Cosío, et al., 2001). As a result, in the last years different electroanatomical mapping systems using different ablation catheters have been increasingly developed to definitively eliminate complex refractory tachyarrhythmias in many patients with or without structural heart diseases, (Nakagawa, Shah, et al., 2001; Dixit, & Callans, 2002; Friedman, 2002; Friedman, Asirvatham, et al., 2002; Kirchhof, Loh, et al., 2002; de Groot, Schalij, et al., 2003; Kopelman, Prater, et al., 2003; Macle, Jais, et al., 2003; Marrouche, Martin, et al., 2003; Oral, Scharf, et al., 2003; Pappone, Rosanio, et al., 2003; Schenider, Ndrepepa, et al., 2003; Pappone, Manguso, et al., 2004; Pappone, Santinelli, et al., 2004; Sporton, Earley, 2004; Ventura, Rostock, et al., 2004; Corrado, Basso, et al., 2005; Gurevitz, Glikson, et al., 2005; Packer, 2005; Rotter, Takahashi, et al., 2005; Earley, Showkathali, et al., 2006; Estner, Deisenhofer, et al., 2006; Kistler, Rajappan, et al., 2006; Oral, Pappone, et al., 2006; Pappone, Augello, et al., 2006; Pappone, & Santinelli, 2006; Roberts-Thomson, Kistler, et al., 2006; Chae, Oral, et al., 2007; Jais, Matsuo, et al., 2009; Martinek, Nesser, et al., 2007; Matsuo, Lim, et al., 2007; Medi, Kalman, et al., 2009; Del Carpio Munoz, Buescher, et al., 2010; Wilber, Pappone, et al., 2010; Khaykin, Oosthuizen, et al., 2011; Barbhuiya, Kumar, et al., 2014; Kang, Etheridge et al., 2014; Kapa, & Asirvatham, 2014; Sun, & Piccini, 2014; Anter, Tschabrunn, & Josephson, 2015; Buch, & Shivkumar, 2015; Ceresnak, Nappo, & Janson, 2015; Scherr, Derval, et al., 2015; N Wang, Ouyang, et al., 2015; Winterfield, Jensen, et al., 2015; Zipes, Calkins, et al., 2015). Since 1990s catheter ablation has been commonly used to treat supraventricular tachyarrhythmias with a success rate of over 90 percent, a low risk of complications while the patient can resume the normal activities in a few days (Silka, Gillette, et al., 1985; Van Hare, Velvis, Langberg, 1990; Kay, Chong, et al., 1993; Kopelman, Prater, et al., 2003; Chen, Chiang, et al., 1994; Lesh, Van Hare, et al., 1994; Poty, 1996; Kopelman, Prater, et al., 2003). It causes little or no discomfort and is done under mild sedation with local anesthesia. In structural heart diseases, catheter ablation is usually performed for drug inefficacy or intolerance or as adjunctive therapy in patients with an implantable cardioverter-defibrillator, who are experiencing frequent DC shock discharges. Consensus groups from the European Society of Cardiology and the North American Society of Pacing and Electrophysiology have defined regular atrial tachycardia (AT) as either focal or macro-reentrant, classified according to electrophysiological mechanisms and anatomical structures (Saoudi, Cosío, et al., 2001). Focal ATs are characterised by a focal origin with subsequent centrifugal spread, with a mechanism based in abnormal automaticity, triggered activity or micro re-entry (Chen, Tai, et al., 1998; Roberts-Thomson, Kistler, et al., 2006). Macro-reentrant ATs, by contrast, involve a larger re-entry circuit, conventionally defined by a diameter in excess of 2 cm. The most common form of macro-reentrant AT involves the cavo-tricuspid isthmus (CTI) as a critical region and has traditionally been referred to as ‘typical atrial flutter’. Since this classification, a third category of atrial tachycardia has been defined. This is termed small circuit re-entry and is “somewhere between micro and macro”. These circuits most commonly occur after atrial fibrillation (AF) ablation procedures involving extensive left atrial ablation. Radiofrequency catheter ablation was first described in pediatric patients in the early 1990s (van Hare, Velvis, & Langberg, 1990). Since then, multiple advances in the technology and understanding of this procedure have allowed this technique to blossom into one of the

26 more pages are available in the full version of this document, which may be purchased using the "Add to Cart" button on the publisher's webpage:  
[www.igi-global.com/chapter/new-imaging-and-computational-technology-as-a-guide-for-catheter-ablation-of-incessant-tachyarrhythmias/149394](http://www.igi-global.com/chapter/new-imaging-and-computational-technology-as-a-guide-for-catheter-ablation-of-incessant-tachyarrhythmias/149394)

## Related Content

---

### A Contactless Fingerprint Verification Method using a Minutiae Matching Technique

Tahirou Djara, Marc Kokou Assogbaand Antoine Vianou (2016). *International Journal of Computer Vision and Image Processing* (pp. 12-27).

[www.irma-international.org/article/a-contactless-fingerprint-verification-method-using-a-minutiae-matching-technique/170593](http://www.irma-international.org/article/a-contactless-fingerprint-verification-method-using-a-minutiae-matching-technique/170593)

### 3D Interaction with Scientific Data Through Virtual Reality and Tangible Interfacing

Wen Qi, Russell M. Taylor, Christopher Healeyand Jean-Bernard Martens (2008). *User Centered Design for Medical Visualization* (pp. 136-173).

[www.irma-international.org/chapter/interaction-scientific-data-through-virtual/30628](http://www.irma-international.org/chapter/interaction-scientific-data-through-virtual/30628)

### Screening of Radiological Images Suspected of Containing Lung Nodules

Raúl Pedro Aceñero Eixarch, Raúl Díaz-Usechi Laplazaand Rafael Berlanga Llavori (2022). *International Journal of Computer Vision and Image Processing* (pp. 1-12).

[www.irma-international.org/article/screening-of-radiological-images-suspected-of-containing-lung-nodules/283968](http://www.irma-international.org/article/screening-of-radiological-images-suspected-of-containing-lung-nodules/283968)

### Translated Trademarks Retrieval using Color Autocorrelogram for Extracted Textual Parts

Ahmed Zeggariand Fella Hachouf (2018). *International Journal of Computer Vision and Image Processing* (pp. 59-67).

[www.irma-international.org/article/translated-trademarks-retrieval-using-color-autocorrelogram-for-extracted-textual-parts/208178](http://www.irma-international.org/article/translated-trademarks-retrieval-using-color-autocorrelogram-for-extracted-textual-parts/208178)

### Effective Video Shot Boundary Detection and Keyframe Selection using Soft Computing Techniques

Rashmi B Sand Nagendraswamy H S (2018). *International Journal of Computer Vision and Image Processing* (pp. 27-48).

[www.irma-international.org/article/effective-video-shot-boundary-detection-and-keyframe-selection-using-soft-computing-techniques/208176](http://www.irma-international.org/article/effective-video-shot-boundary-detection-and-keyframe-selection-using-soft-computing-techniques/208176)