Chapter 36 QSPR/QSAR Analyses by Means of the CORAL Software: Results, Challenges, Perspectives

Andrey A. Toropov IRCCS-Istituto di Ricerche Farmacologiche Mario Negri, Italy

Alla P. Toropova IRCCS-Istituto di Ricerche Farmacologiche Mario Negri, Italy

Emilio Benfenati IRCCS-Istituto di Ricerche Farmacologiche Mario Negri, Italy

Orazio Nicolotti Università degli Studi di Bari "Aldo Moro", Italy

Angelo Carotti Università degli Studi di Bari "Aldo Moro", Italy

Karel Nesmerak *Charles University in Prague, Czech Republic*

> Aleksandar M. Veselinović University of Niš, Serbia

Jovana B. Veselinović University of Niš, Serbia

Pablo R. Duchowicz Instituto de Investigaciones Fisicoquímicas Teóricas y Aplicadas INIFTA (UNLP, CCT La Plata-CONICET), Argentina

> **Daniel Bacelo** Universidad de Belgrano, Argentina

Eduardo A. Castro Instituto de Investigaciones Fisicoquímicas Teóricas y Aplicadas INIFTA (UNLP, CCT La Plata-CONICET), Argentina

> Bakhtiyor F. Rasulev Jackson State University, USA

> **Danuta Leszczynska** Jackson State University, USA

> Jerzy Leszczynski Jackson State University, USA

ABSTRACT

In this chapter, the methodology of building up quantitative structure—property/activity relationships (QSPRs/QSARs)—by means of the CORAL software is described. The Monte Carlo method is the basis of this approach. Simplified Molecular Input-Line Entry System (SMILES) is used as the representation of the molecular structure. The conversion of SMILES into the molecular graph is available for QSPR/QSAR analysis using the CORAL software. The model for an endpoint is a mathematical function of

DOI: 10.4018/978-1-5225-1762-7.ch036

the correlation weights for various features of the molecular structure. Hybrid models that are based on features extracted from both SMILES and a graph also can be built up by the CORAL software. The conceptually new ideas collected and revealed through the CORAL software are: (1) any QSPR/QSAR model is a random event; and (2) optimal descriptor can be a translator of eclectic information into an endpoint prediction.

INTRODUCTION

Almost seventy years ago one of the areas of theoretical chemistry named the quantitative structure – property / activity relationships (QSPRs/QSARs) was established (Wiener, 1947, pp. 17-20; Wiener, 1948, pp. 425-430). However, if one considers the publication of Hammett's Free-energy relationship (Hammett, 1935, pp. 125–136) as the first exercise in the QSPR/QSAR analyses, then the introduction of this area of theoretical chemistry should be moved back by eighty years. This is obviously an evidence of the long lasting importance of the QSAR approaches. Such significance is due to their wide applications. The main target of QSPR/QSAR analyses is the prediction of the numerical data related to various endpoints (physicochemical in the case of QSPR; and biochemical, ecological, medicinal, etc., in the case of QSAR).

One can consider three periods of evolution in this field.

- The first period can be defined as the construction of molecular descriptors and establishing of their correlation with various endpoints (Bonchev & Trinajstic, 1977; Balaban, 1982, pp. 399-404; Bonchev, 1995, pp. 137- 156; Diudea & Gutman, 1998; Ivanciuc, Ivanciuc, & Balaban, 1998; Randic, DeAlba, & Harris, 1998; Castro, Tueros, & Toropov, 2000; Randic & Basak, 2001; Nicolic, Kovacevic, Milicevic, & Trinajstic, 2003; Ren, 2003, pp. 29-39; González, Terán, Teijeira, & González-Moa, 2005; Melagraki, Afantitis, Sarimveis, Igglessi-Markopouloua, & Supuran, 2006).
- 2. The second period can be defined as the construction of molecular descriptors, establishing of their correlation with an endpoint for the compounds of "visible" training set with the further testing of the predictive potential of this correlation with "invisible" compounds (i.e. compounds which are not involved in building up of the model) involved in the test set (Cronin et al., 2002; Golbraikh & Tropsha, 2002; Taskinen & Yliruusi, 2003; Hemmateenejad, 2004, pp. 475–485; Duchowicz, Castro, Fernandez, & Gonzalez, 2005; Oberg, 2005, pp. 2189–2200; Afantitis et al., 2006; Coi et al., 2006; Leonard & Roy, 2006; Pan, Jiang, & Wang, 2007; Roy & Roy, 2008; Porto, Souza, Junkes, Yunes, & Heinzen, 2008; Toropova, Toropova, Benfenati, Leszczynska, & Leszczynski, 2010; Toropov et al., 2012a; Toropov, Toropova, Raska Jr, Benfenati, & Gini, 2012b; Nesmerak, Toropov, Toropova, & Kohoutova, 2013; Papa, van derWal, Arnot, & Gramatica, 2014).
- The third period can be defined as building up of the QSPR/QSAR models according to OECD principles (Kruhlak, Contrera, Daniel Benz, & Matthews, 2007; Kar & Roy, 2010, 2012; Putz, Ionascu, Putz, & Ostafe, 2011; de Melo, 2012, pp. 213–222; Sahigara et al., 2012; Toropova & Toropov, 2014).

According to the OECD principles, the QSPR/QSAR model should be characterized by:

25 more pages are available in the full version of this document, which may be purchased using the "Add to Cart" button on the publisher's webpage: www.igi-global.com/chapter/qsprqsar-analyses-by-means-of-the-coralsoftware/174156

Related Content

A Study of Third-Semester Traditional Chinese Medicine Science Students

Yinuo Wang (2023). Cases on Teaching Pharmacology to Complementary and Alternative Medicine Students (pp. 38-52).

www.irma-international.org/chapter/a-study-of-third-semester-traditional-chinese-medicine-science-students/320630

Resealed Erythrocytes as Drug Carriers and Its Therapeutic Applications

Prabhakar Singh, Sudhakar Singhand Rajesh Kumar Kesharwani (2017). *Pharmaceutical Sciences: Breakthroughs in Research and Practice (pp. 459-485).* www.irma-international.org/chapter/resealed-erythrocytes-as-drug-carriers-and-its-therapeutic-applications/174136

Antioxidant Activity and Phytochemical Composition of the Selected Fruits

D. Jancy Rani (2024). *Ethnobotanical Insights Into Medicinal Plants (pp. 1-19).* www.irma-international.org/chapter/antioxidant-activity-and-phytochemical-composition-of-the-selected-fruits/346938

Marine Bioactive Products as Potential Antileishmanial Therapeutics

Muhammad Adil, Pragya Tiwari, Mavara Iqbal, Shamsa Kanwaland Aisha Umar (2023). *Pharmacological Benefits of Natural Agents (pp. 225-249).*

www.irma-international.org/chapter/marine-bioactive-products-as-potential-antileishmanial-therapeutics/327311

Ethnobotanical and Pharmacological Importance of the Herbal Plants With Anti-Hair Fall and Hair Growth Activities

Kiran Mustafa, Javaria Kanwal, Sara Musaddiqand Samia Khakwani (2020). Advanced Pharmacological Uses of Medicinal Plants and Natural Products (pp. 49-66).

www.irma-international.org/chapter/ethnobotanical-and-pharmacological-importance-of-the-herbal-plants-with-anti-hair-fall-and-hair-growth-activities/252935