

# Chapter 4

## Store–Operated Calcium Entry Channels: Potential Role in Cardiac Function

**Diptiman D. Bose**  
Western New England University, USA

### ABSTRACT

*Store-operated  $\text{Ca}^{2+}$  entry (SOCE) channels mediate  $\text{Ca}^{2+}$  influx from the extracellular milieu into the cytosol to regulate a myriad of cellular functions. The  $\text{Ca}^{2+}$ -release activated  $\text{Ca}^{2+}$  current has been well characterized in non-excitabile cells such as immune cells. However, the role of SOCE proteins in cardiomyocytes and cardiac function has only been recently investigated. The localized endoplasmic reticulum protein, stromal interaction molecule (STIM) and plasma membrane  $\text{Ca}^{2+}$  channels, ORAI form the minimal functional unit of SOCE. The documentation of STIM and Orai expression in cardiomyocytes has raised questions regarding their role in cardiac function. Recent evidence supports the central role of STIM and Orai in gene transcription and, subsequent phenotypic changes associated with cardiac remodeling and hypertrophy. The purpose of this chapter is to provide an overview of our current understanding of SOCE proteins and, to explore their contributions to cardiovascular function and role in cardiac disorders.*

### INTRODUCTION

Calcium's role as a ubiquitous intracellular messenger is demonstrated by its central role in a wide range of cellular functions from cell growth, proliferation, function and even cell death. In order to elicit cellular response, a cell recruits various pumps, exchangers and channels to regulate the concentration of  $\text{Ca}^{2+}$  (Berridge, Bootman, & Roderick, 2003; Prakriya & Lewis, 2015; J. W. Putney, 2011). The endoplasmic reticulum/sarcoplasmic reticulum (ER/SR) act as an intracellular storage of  $\text{Ca}^{2+}$ , while plasma membrane (PM) channels regulate gating of  $\text{Ca}^{2+}$  from the extracellular space. Store operated  $\text{Ca}^{2+}$  entry (SOCE), is a major mechanism representing  $\text{Ca}^{2+}$  entry in many excitable and non-excitabile cells. Since the first characterization of SOCE through electrophysiology, the two major components of the SOCE

DOI: 10.4018/978-1-5225-2092-4.ch004

process were identified; the ER/SR  $\text{Ca}^{2+}$  sensor, stromal interaction molecule (STIM) and, the plasma membrane (PM) localized Orai protein as the store-operated channels (Fahrner, Derler, Jardin, & Romanin, 2013; J. W. Putney, 2011; Shaw & Feske, 2012). Upon depletion of inositol-1, 4, 5-trisphosphate ( $\text{IP}_3$ ) or Ryanodine (RyR)-sensitive ER/SR stores, localized ER sensor STIM directly couples with PM Orai channels mediating  $\text{Ca}^{2+}$  influx. Since the depletion of ER/SR  $\text{Ca}^{2+}$  is the trigger for PM  $\text{Ca}^{2+}$  entry, this pathway was appropriately named “store operated  $\text{Ca}^{2+}$  entry”. SOCE-mediated  $\text{Ca}^{2+}$  entry not only allows for refilling of the ER/SR stores but also helps maintain  $\text{Ca}^{2+}$  homeostasis. The sustained entry of  $\text{Ca}^{2+}$  also serves other purposes such as activation of secretion, modulation of enzyme activation, and initiation of transcriptional signaling (J. W. Putney, 2011). Since SOCE-mediated  $\text{Ca}^{2+}$  influx is involved in vital cellular processes, it is not surprising that aberrant SOCE function has been implicated in many disease states including immunodeficiency, acute pancreatitis, Alzheimer’s disease, Duchenne muscular dystrophy and cardiac hypertrophy (Karlstad, Sun, & Singh, 2012; J. W. Putney, 2011).

Since its discovery, the SOCE phenomenon has been well characterized in non-excitable immune cells (Prakriya & Lewis, 2015; Shaw & Feske, 2012). Soon after, SOCE-induced  $\text{Ca}^{2+}$  influx was shown to play a critical role in excitable cell such as neurons, skeletal muscle cells, and cardiomyocytes (Hartmann et al., 2014; Liu, Xin, Benson, Allen, & Ju, 2015; Majewski & Kuznicki, 2015; Stiber et al., 2008; Tojyo, Morita, Nezu, & Tanimura, 2014). Several studies have demonstrated the expression of STIM and Orai in adult cardiomyocytes, ventricular myocardium, and the sinoatrial node (Wolkowicz et al., 2011; Zhu-Mauldin, Marsh, Zou, Marchase, & Chatham, 2012). In fact, strong evidence suggests that the STIM1 and Orai play a key role in the progression of cardiac hypertrophy. Recent investigations have shown the increased expression of STIM1 in a hypertrophic response (Collins, Zhu-Mauldin, Marchase, & Chatham, 2013). With advances in molecular techniques and transgenic models, studies have provided insight into the role of Orai and transient receptor potential (TRP) channels in the etiology of several cardiovascular diseases (Yue et al., 2015). The goal of this chapter is to provide an overview of our current understanding of molecular regulation of SOCE and highlight the role of STIM1/Orai-1-mediated SOCE in cardiomyocyte function and pathology.

## **STORE OPERATED CALCIUM ENTRY**

The ligation of agonists such as growth factors, neurohormonal stimuli, and inflammatory mediators to G-protein coupled receptors or receptor tyrosine kinases initiates the activation of phospholipase (PLC) enzymes. PLC hydrolyzes phosphatidylinositol 4, 5 biphosphate into diacylglycerol and inositol 1, 4, 5-trisphosphate ( $\text{IP}_3$ ).  $\text{IP}_3$  binds to the  $\text{IP}_3$  receptor ( $\text{IP}_3\text{R}$ ) on the ER/SR, triggering  $\text{Ca}^{2+}$  release into the cytosol and mobilizing an increase in cytosolic  $\text{Ca}^{2+}$  ( $[\text{Ca}^{2+}]_i$ ) (Berridge et al., 2003; Oh-hora & Rao, 2008). The increase in  $[\text{Ca}^{2+}]_i$  can fuel activation of localized events such as calcium-induced  $\text{Ca}^{2+}$  release via the Ryanodine receptors (RyRs) or even induce downstream transcriptional events (Bers, 2008; J. W. Putney, 2011). The discovery of  $\text{IP}_3$  led to our understanding that the consequence of  $\text{IP}_3$  production was a transient increase in  $[\text{Ca}^{2+}]_i$ , followed by a sustained influx of  $\text{Ca}^{2+}$  from the extracellular space. While the increase in  $[\text{Ca}^{2+}]_i$  was identified as a product of  $\text{IP}_3$  activating the  $\text{IP}_3\text{R}$  in the ER/SR, the mechanism underlying  $\text{Ca}^{2+}$  influx remained unclear (Berridge, 1993). Several studies suggested that a PM- $\text{Ca}^{2+}$  pathway was responsible for directly reloading the ER/SR stores. The idea of SOCE, initially described as “capacitative  $\text{Ca}^{2+}$  entry (CCE)”, first described by Putney, reflected these conclusions. This model suggested that  $\text{Ca}^{2+}$  from extracellular space directly loaded the ER (the “capacitor”) (J.

18 more pages are available in the full version of this document, which may be purchased using the "Add to Cart" button on the publisher's webpage:  
[www.igi-global.com/chapter/store-operated-calcium-entry-channels/176209](http://www.igi-global.com/chapter/store-operated-calcium-entry-channels/176209)

## Related Content

---

### Interdisciplinary Approach to Cardiovascular Diseases for Research and Everyday Clinical Practice Purposes

Aleksander Goch, Anna Rosiek, Krzysztof Leksowski and Emilia Mikoajewska (2019). *Coronary and Cardiothoracic Critical Care: Breakthroughs in Research and Practice* (pp. 139-171).  
[www.irma-international.org/chapter/interdisciplinary-approach-to-cardiovascular-diseases-for-research-and-everyday-clinical-practice-purposes/225360](http://www.irma-international.org/chapter/interdisciplinary-approach-to-cardiovascular-diseases-for-research-and-everyday-clinical-practice-purposes/225360)

### Store-Operated Calcium Entry Channels: Potential Role in Cardiac Function

Diptiman D. Bose (2017). *Emerging Applications, Perspectives, and Discoveries in Cardiovascular Research* (pp. 53-72).  
[www.irma-international.org/chapter/store-operated-calcium-entry-channels/176209](http://www.irma-international.org/chapter/store-operated-calcium-entry-channels/176209)

### Coronary Artery Disease

Rakesh C. Arora and Rohit Singal (2015). *Modern Concepts and Practices in Cardiothoracic Critical Care* (pp. 643-682).  
[www.irma-international.org/chapter/coronary-artery-disease/136926](http://www.irma-international.org/chapter/coronary-artery-disease/136926)

### Thoracic Aortic Disease

Berhane Worku and Leonard Girardi (2015). *Modern Concepts and Practices in Cardiothoracic Critical Care* (pp. 728-755).  
[www.irma-international.org/chapter/thoracic-aortic-disease/136929](http://www.irma-international.org/chapter/thoracic-aortic-disease/136929)

### Wearable Technology for Cardiology

(2023). *Wearable and Implantable Electrocardiography for Early Detection of Cardiovascular Diseases* (pp. 96-108).  
[www.irma-international.org/chapter/wearable-technology-for-cardiology/329312](http://www.irma-international.org/chapter/wearable-technology-for-cardiology/329312)