

# Chapter 10

## A Transaction Cost Economics Perspective for Pervasive Technology

**Nilmini Wickramasinghe**

*Epworth HealthCare, Australia & Deakin University, Australia*

**Indrit Troshani**

*University of Adelaide, Australia*

**Steve Goldberg**

*INET International Inc., Canada*

### ABSTRACT

*Numerous mobile technology solutions are being developed and implemented today to address a myriad of healthcare problems. However, it remains unclear what the true cost/benefit of these solutions is and who benefits from them. To investigate this we apply a transaction cost economics framework to a pervasive mobile solution that has been designed and developed to enhance diabetes self-care. Diabetes is one of the leading chronic diseases and its prevalence continues to rise. The solution examined in this paper relies on pervasive wireless technology and is designed to facilitate the effective management of diabetes in the context of gestational diabetes, a conditions that affects up to 8% of pregnant women. A transactions cost assessment of this solution is provided.*

### INTRODUCTION

Mobile technology has created many opportunities for a plethora of new applications to emerge. More recently, many of these new applications are focused on addressing health and wellness issues such as obesity, diabetes, and wellness aspects including nutritional intake, and exercise. Many believe that these applications will assist in addressing the current health concerns including diabetes and obesity; however it is still unclear how cost effective these solutions really are and who benefits from them. In

DOI: 10.4018/978-1-5225-3926-1.ch010

an attempt to shed light on these issues and guide existing and future research, we examine a specific pervasive mobile solution that is designed to enable diabetes self-care. We assess this solution the transaction economics cost perspective to assess its key cost and benefit aspects. The study contributes to both theory and practice.

## **BACKGROUND**

Diabetes mellitus is one of the leading chronic diseases affecting Australians and its prevalence continues to rise exponentially. The total number of diabetes patients worldwide is estimated to rise to 366 million in 2030 from 171 million in 2000 (Wild, Roglic, Green, Sicree, & King, 2004). With increasingly growing prevalence which includes an estimated 275 Australians developing diabetes daily (DiabetesAustralia, 2008), Australia is expected to be a significant contributor to this projected trend. An estimated 700,000 Australians, representing approximately 3.6% of the population, were diagnosed with diabetes in 2004-05. Between 1989-90 and 2004-05 the proportion of Australians diagnosed with this disease more than doubled from 1.3% to 3.3%. Additionally, between 2000-01 and 2004-05, Australian diabetes hospitalizations increased by 35% from 1,932 to 2,608 hospitalizations per 100,000 people (AIHW, 2007, 2008). For every person diagnosed with diabetes, it is estimated that there is another who has yet to be diagnosed, which doubles the number of diabetes sufferers (DiabetesAustralia, 2008). Diabetes is, thus, one of the fastest growing chronic diseases in Australia (AIHW, 2008; Catanzariti, Faulks, & Waters, 2007; Chittleborough, Grant, Phillips, & Taylor, 2007). Diabetes and its complications incur significant costs for the health system in Australia, including costs incurred by carers, government, and the entire health system (DiabCostAustralia, 2002). In 2004-05 direct healthcare expenditure on diabetes was A\$907 million, which constituted approximately 2% of the allocatable recurrent health expenditure in that year (AIHW, 2007, 2008). Further costs include societal costs that represent productivity losses for both patients and their carers (DiabCostAustralia, 2002).

Gestational diabetes mellitus (GDM) is a common form of diabetes that presents in pregnancy, sometimes with symptoms but often diagnosed in otherwise normal women on routine screening tests. Some women, particularly those in whom the diagnosis of GDM is made early in pregnancy, may have pre-existing undiagnosed diabetes. In Australia and New Zealand, universal screening for GDM is recommended by the Australasian Diabetes in Pregnancy Society (ADIPS) (Hoffman, Nola, Wilson, Oats, & Simmons, 1998), although the uptake of this recommendation is variable (Rumbold & Crowther, 2001). It is estimated that in Australia, 3-6% of pregnant women will develop GDM at around 24-28 weeks gestation, with a smaller number earlier and later in pregnancy. An Australian study of 210 pregnant women found that screening for GDM had an adverse impact on women's perceptions of their own health (Rumbold & Crowther, 2001, 2002). GDM is more common in older women, in those with a family history of diabetes, in those who are overweight, and in those of non-Caucasian heritage (Carolan, Steele, & Margetts, 2010). Maternal complications of GDM can be serious and include polyhydramnios and premature labour, maternal hypertension, low birth weights and stillbirth (Fan, Yang, Gao, Lintu, & Sun, 2006; Hoffman et al., 1998). It recurs in subsequent pregnancy in 30-80% of women, the incidence varying with ethnicity, being lower in Caucasian women (Kim, Berger, & Chamany, 2007).

Treatment of women with GDM aims to control maternal, and therefore fetal, hyperglycaemia and the associated tendency of fetal hyperinsulinaemia which is at the root of the fetal complications (Metzger &

21 more pages are available in the full version of this document, which may be purchased using the "Add to Cart" button on the publisher's webpage:

[www.igi-global.com/chapter/a-transaction-cost-economics-perspective-for-pervasive-technology/192673](http://www.igi-global.com/chapter/a-transaction-cost-economics-perspective-for-pervasive-technology/192673)

## Related Content

---

### A Prototype Agent Based Model and Machine Learning Hybrid System for Healthcare Decision Support

Marek Laskowski (2011). *International Journal of E-Health and Medical Communications* (pp. 67-90).

[www.irma-international.org/article/prototype-agent-based-model-machine/60207](http://www.irma-international.org/article/prototype-agent-based-model-machine/60207)

### A Multi-Agent Based Modeling and Simulation Data Management and Analysis System for the Hospital Emergency Department

Manel Saad Saoud, Abdelhak Boubetra and Safa Attia (2017). *International Journal of Healthcare Information Systems and Informatics* (pp. 21-36).

[www.irma-international.org/article/a-multi-agent-based-modeling-and-simulation-data-management-and-analysis-system-for-the-hospital-emergency-department/182480](http://www.irma-international.org/article/a-multi-agent-based-modeling-and-simulation-data-management-and-analysis-system-for-the-hospital-emergency-department/182480)

### A Threat Table Based Assessment of Information Security in Telemedicine

John C. Pendergrass, Karen Heart, C. Ranganathan and V. N. Venkatakrishnan (2014). *International Journal of Healthcare Information Systems and Informatics* (pp. 20-31).

[www.irma-international.org/article/a-threat-table-based-assessment-of-information-security-in-telemedicine/124117](http://www.irma-international.org/article/a-threat-table-based-assessment-of-information-security-in-telemedicine/124117)

### Planning and Control and the Use of Information Technology in Mental Healthcare Organizations

I.J. Baars and G.G. Van Merode (2010). *Health Information Systems: Concepts, Methodologies, Tools, and Applications* (pp. 1066-1073).

[www.irma-international.org/chapter/planning-control-use-information-technology/49917](http://www.irma-international.org/chapter/planning-control-use-information-technology/49917)

### RMAC: Customised MAC Protocol for Roundabout Management Using VANET for Cooperative Driving

Hitender Vats and Ranjeet Singh Tomar (2021). *International Journal of E-Health and Medical Communications* (pp. 77-92).

[www.irma-international.org/article/rmac/267956](http://www.irma-international.org/article/rmac/267956)