

Chapter 26

Use of GIS and Remote Sensing for Landslide Susceptibility Mapping

Arzu Erener

Kocaeli University, Turkey

Gulcan Sarp

Suleyman Demirel University, Turkey

Sebnem Duzgun

Middle East Technical University, Turkey

ABSTRACT

In recent years, geographical information systems (GISs) and remote sensing (RS) have proven to be common tools adopted for different studies in different scientific disciplines. GIS is defined as a set of tools for the input, storage, retrieval, manipulation, management, modeling, analysis, and output of spatial data. RS, on the other hand, can play a role in the production of a data and in the generation of thematic maps related to spatial studies. This study focuses on use of GIS and RS data for landslide susceptibility mapping. Five factors including normalized difference vegetation index (NDVI) and topographic wetness index (TWI), slope, lineament density, and distance to roads were used for the grid-based approach for landslide susceptibility mappings. Results of this study suggest that geographic information systems can effectively be used to obtain susceptibility maps by compiling and overlaying several data layers relevant to landslide hazards.

INTRODUCTION

In recent years, geographical information systems (GISs) and Remote Sensing (RS) have proven to be common tools adopted for various studies in different scientific disciplines. GIS provides as a set of tools for the input, storage, retrieval, manipulation, management, modeling, analysis and output of spatial data. RS, on the other hand, offers earth observation data for thematic maps related to spatial studies. The use

DOI: 10.4018/978-1-5225-7359-3.ch026

Use of GIS and Remote Sensing for Landslide Susceptibility Mapping

of GIS and RS data for landslide susceptibility mapping are demonstrated by three different landslide susceptibility maps with five different variables (Normalized Difference Vegetation Index (NDVI), Topographic Wetness Index (TWI), slope; lineament density and distance to roads). The comparison of the generated final susceptibility maps with historical landslide locations is given with important factors affecting the accuracy of susceptibility map. The accuracy analysis of the final susceptibility maps for various weighting strategies is performed. The results indicate that assignment of weights to the slope parameter impacts the accuracy in the high susceptible zones.

BACKGROUND

Landslides are among the most common natural hazards and are the most damaging, leading to substantial economic, human, and environmental losses throughout the world. The quantitative assessment of landslide hazards for a large area is critical for mitigation of the associated risks. They are often triggered by natural phenomena and/or human activity, such as earthquakes, precipitation, erosion, deforestation etc. and are difficult to predict. One of the greatest limiting factors in predicting and mapping landslide activity is the lack of understanding of scale-dependent processes, such as erosion, weathering, and fracturing (Glenn et al., 2006). Such maps normally aim at providing a document that depicts the likelihood or possibility of new movements occurring in an area, and therefore helping to reduce future damages. To express the potential for occurrence of landslides in a quantitative manner, maps must incorporate the concept of probability, which is an assessment of the relative frequency of occurrence (Ohlmacher & Davis, 2003). Susceptibility expresses the likelihood that a landslide will occur in an area on the basis of the local terrain conditions (Soeters & Van Westen, 1996); return period or annual probability of occurrence is not considered. The main difference between susceptibility and hazard is therefore that the latter considers the temporal factor, by estimating the probability of occurrence of the phenomenon within a specified period of time (Varnes, 1984) whereas the former considers the likelihood of landslide occurrence. There are many studies in the literature about use of GIS to evaluate landslide susceptibility (e.g. Gokceoglu et al. 2005; Akgun et al. 2011; Akgun 2012; Pradhan et al. 2013; Kavzoglu et al. 2013)

MAIN FOCUS

The purpose of this study is to apply the grid based GIS techniques for landslide susceptibility mapping using five different factors including Normalized Difference Vegetation Index (NDVI), Topographic Wetness Index (TWI), slope, lineament density, and distance to roads. The scope includes the preparation of landslide susceptibility map to identify highly susceptible areas and, the accuracy assessment related to the obtained maps.

The susceptibility assessment methodology is demonstrated for More and Romsdal region in Norway (Erener & Duzgun, 2010). The study area occupies approximately 606.755 km² in the west part of Norway. The upper left coordinates on 112707,770408 m - 6952112,603469 m and lower right coordinates 6929466,479194 m -144909,272731 m respectively (Figure 1).

13 more pages are available in the full version of this document, which may be purchased using the "Add to Cart" button on the publisher's webpage:

www.igi-global.com/chapter/use-of-gis-and-remote-sensing-for-landslide-susceptibility-mapping/211885

Related Content

Role of Plant Growth Promoting Rhizosphere (PGPR) on Molecular Mechanisms Transporters Under Heavy Metal Stress

(2020). *Nano-Phytoremediation Technologies for Groundwater Contaminates: Emerging Research and Opportunities* (pp. 71-84).

www.irma-international.org/chapter/role-of-plant-growth-promoting-rhizosphere-pgpr-on-molecular-mechanisms-transporters-under-heavy-metal-stress/241169

Indian National Strategy for Climate Change Adaptation and Mitigation

B. K. Khanna (2018). *Climate Change and Environmental Concerns: Breakthroughs in Research and Practice* (pp. 541-572).

www.irma-international.org/chapter/indian-national-strategy-for-climate-change-adaptation-and-mitigation/201722

Measuring Dynamics of Ecological Footprint as an Index of Environmental Sustainability at the Regional Level Using Geospatial Information Technology: Measuring Ecological Footprint Using GIS

Laxmikant Sharma and Suman Sinha (2019). *Environmental Information Systems: Concepts, Methodologies, Tools, and Applications* (pp. 965-980).

www.irma-international.org/chapter/measuring-dynamics-of-ecological-footprint-as-an-index-of-environmental-sustainability-at-the-regional-level-using-geospatial-information-technology/212977

Impact of Global Climate Change on Potato Diseases and Strategies for Their Mitigation

Mehi Lal, Saurabh Yadav, Rajendra Prasad Pant, Vijay Kumar Dua, B. P. Singh and Surinder Kumar Kaushik (2018). *Climate Change and Environmental Concerns: Breakthroughs in Research and Practice* (pp. 134-151).

www.irma-international.org/chapter/impact-of-global-climate-change-on-potato-diseases-and-strategies-for-their-mitigation/201697

Cold Thermal Energy Storage

Franc Franc Kosi, Branislav Zivkovic, Mirko S. Komatina, Dragi Antonijevic, Mohamed Abdul Galil and Uros Milorad Milovancevic (2017). *Renewable and Alternative Energy: Concepts, Methodologies, Tools, and Applications* (pp. 93-123).

www.irma-international.org/chapter/cold-thermal-energy-storage/169593