

## Chapter 43

# Monitoring Changes in Urban Cover Using Landsat Satellite Images and Demographical Information

**Prashant K. Srivastava**

*Banaras Hindu University, India*

**Swati Suman**

*Banaras Hindu University, India*

**Smita Pandey**

*Banaras Hindu University, India*

### ABSTRACT

*The monitoring of urban cover is very important for the planner, management, governmental and non-governmental organizations for optimizing the use of urban resources and minimizing the environmental losses. The study here aims at analyzing the changes that occurred in urban green cover over a time span of 1991-2001 using multi-date Landsat satellite images data over the Varanasi district, India and its relation to demographical changes. The Support Vector Machines (SVMs) classifier has been used for image classification. The urbanization indicators such as Land Consumption Ratio (LCR) and Land Absorption Coefficient (LAC) were also used in order to understand the changes in urban cover and population dynamics. All the analysis indicates significant changes in the urban cover values with increasing population at both spatial and temporal scale.*

### INTRODUCTION

Technological advancement in satellite remote sensing leads to a substantial development in the environmental and resource management policies in recent decades. Satellite earth imagery due to their global coverage and continuous temporal monitoring are proved to be significant in today's chaotic scenario of expedited population growth for limited natural resources over traditional practices (Srivastava et al.,

DOI: 10.4018/978-1-5225-7033-2.ch043

2010). Developmental strategies in alliance with land use land cover change monitoring have laid to a new prospect towards sustainable development especially for a mercurially evolving country like India (Singh et al., 2016).

Land cover is the state earth surface and its biophysical attributes (Lambin et al., 2001), whereas land use is the manner in which humans use natural resources for their use and convenience (Srivastava et al., 2012). Land use land cover change is significant to a range of themes to study and simulate global changes as land covers are a subject of change with pace, magnitude and spatial reach of human beings suggested by leading experts (Singh et al., 2014). Urbanization, deforestation and desertification are the consequences of uninhibited development policies due to modern territorial and socio-economic progress leading to the transformation of existing land use land cover pattern (Srivastava et al., 2013). Since deforestation and desertification needs immense amount of ecological and societal awareness and efforts to be controlled, a systematic expansion concrete sector can be our evident call for achieving sustainable development goals (Singh et al., 2015).

Urban expansion in developing countries world has been dramatic and rapid. United Nations in a report published in year 2015 quoted that the “total urban population in developing countries like India and china has increased 10 folds from 300 million to 3 billion since 1950”. Coupled with the technological advancements and population explosion, Urbanization has accelerated industrial revolution leading to increased resource consumption and environment degradation rates (Banerjee & Srivastava, 2013). Rapid urbanization has been a topic of interest for researcher, planners economist, ecologist and policy makers as these developmental advances are happening in an era of shooting population, shrinking natural resource and increasing demands in need to ensure sustainable development for all (Srivastava et al., 2014b).

Digital change detection is a process of surveying temporal changes for a place outlined over ecological entities through multi temporal satellite images by detecting changes in radiance value over time (Srivastava et al., 2014a). Change detection exercises have a long range implication in

- Urban sprawl studies (Jat et al., 2008),
- Land use analysis (Martinuzzi et al., 2007),
- Monitoring shifting cultivation (Roy et al., 1985),
- Forest assessment (Wang et al., 2010),
- Deforestation mapping (Frohn et al., 1996),
- Snow melt measurement (Abdalati & Steffen, 1997),
- Disaster monitoring (Dlamini, 2009),
- Damage assessment (Rejaie and Shinozuka, 2004) and many more.

Combined assessment of satellite image classification with demographic status of the area can be used to monitor urban sprawl. Demography of an area significantly influences development and architectural pattern of an area. These studies are often supported through either an image-to-image classification or a post- classification comparison method. The post classification method compares two or more separately classified images of different dates and is considered too be most appropriate and conveniently used approach (Singh, 1989).

13 more pages are available in the full version of this document, which may be purchased using the "Add to Cart" button on the publisher's webpage:

[www.igi-global.com/chapter/monitoring-changes-in-urban-cover-using-landsat-satellite-images-and-demographical-information/212978](http://www.igi-global.com/chapter/monitoring-changes-in-urban-cover-using-landsat-satellite-images-and-demographical-information/212978)

## Related Content

---

### Electronic and ICT Solutions for Smart Buildings and Urban Areas

Luca Tamburini, Maurizio Rossi and Davide Brunelli (2017). *Renewable and Alternative Energy: Concepts, Methodologies, Tools, and Applications* (pp. 1776-1804).

[www.irma-international.org/chapter/electronic-and-ict-solutions-for-smart-buildings-and-urban-areas/169658](http://www.irma-international.org/chapter/electronic-and-ict-solutions-for-smart-buildings-and-urban-areas/169658)

### GIS, Spatial Analysis, and Modeling: The Case of Breast Cancer Incidence in the US

Khadijeh Rouzbehani, Ghazaleh Sajjadi and Mohamad Rahim Hatami (2019). *Environmental Information Systems: Concepts, Methodologies, Tools, and Applications* (pp. 1284-1297).

[www.irma-international.org/chapter/gis-spatial-analysis-and-modeling/212994](http://www.irma-international.org/chapter/gis-spatial-analysis-and-modeling/212994)

### Identification of Geospatial Objects Using Spectral Pattern

Subhabrata Barman (2019). *Environmental Information Systems: Concepts, Methodologies, Tools, and Applications* (pp. 882-893).

[www.irma-international.org/chapter/identification-of-geospatial-objects-using-spectral-pattern/212973](http://www.irma-international.org/chapter/identification-of-geospatial-objects-using-spectral-pattern/212973)

### Adaptations to Climate Change and Climate Variability in the Agriculture Sector in Mauritius: Lessons from a Technical Needs Assessment

Prakash N. K. Deenapanray and Indoomatee Ramma (2015). *Impacts of Climate Change on Food Security in Small Island Developing States* (pp. 130-165).

[www.irma-international.org/chapter/adaptations-to-climate-change-and-climate-variability-in-the-agriculture-sector-in-mauritius/118023](http://www.irma-international.org/chapter/adaptations-to-climate-change-and-climate-variability-in-the-agriculture-sector-in-mauritius/118023)

### Decreasing Wear of Large Vertical Axis Wind Turbines by Employing a Multi-Level Turbine Concept

Jan H. Winiewski (2017). *Renewable and Alternative Energy: Concepts, Methodologies, Tools, and Applications* (pp. 1015-1028).

[www.irma-international.org/chapter/decreasing-wear-of-large-vertical-axis-wind-turbines-by-employing-a-multi-level-turbine-concept/169623](http://www.irma-international.org/chapter/decreasing-wear-of-large-vertical-axis-wind-turbines-by-employing-a-multi-level-turbine-concept/169623)