

Chapter 8

The Importance of Ionic Liquids and Applications on Their Molecular Modeling

Sefa Celik

Istanbul University – Cerrahpasa, Turkey

Ali Tugrul Albayrak

Istanbul University – Cerrahpasa, Turkey

Sevim Akyuz

Istanbul Kultur University, Turkey

Aysen E. Ozel

Istanbul University, Turkey

ABSTRACT

Ionic liquids are salts with melting points generally below 100 °C made of entirely ions by the combination of a large cation and a group of anions. Some ionic liquids are found to have therapeutic properties due to their toxic effects (e.g., anticancer, antibacterial, and antifungal properties). The determination of the most stable molecular structures, that is, the lowest energy conformer of these ionic liquids with versatile biological activities, is of particular importance. Density function theory (DFT) based on quantum mechanical calculation method, one of the molecular modeling methods, is widely used in physics and chemistry to determine the electronic structures of these stable geometries and molecules. With the theory, the energy of the molecule is determined by using the electron density instead of the wave function. It is observed that the theoretical models developed on the ionic liquids in the literature are in agreement with the experimental results because of electron correlations included in the calculation.

DOI: 10.4018/978-1-5225-7467-5.ch008

INTRODUCTION

The first ionic liquid was synthesized by Paul Walden 104 years ago. Ionic liquids are eutectic salt mixtures that consist completely of ions and their melting point is normally below 100 °C. Ionic liquids have been of interest by reason of their many unique features such as superior capacity to dissolve organic and inorganic substances, enormously low volatility, adjustable nature, high thermal stability, high chemical inertness, adjustable viscosity, high conductivity, high heat capacity, wide electrochemical range. Ionic liquids have outstanding handling advantages like nearly zero probability of explosion in high-temperature chemical reactions on account of their low volatilities and non-flammability, their use over a broad range of reaction temperatures owing to their high thermal stabilities (ie thermal decay generally above 350 °C) and high boiling points, dissolution of several organic and inorganic materials in this liquids and usage of polar aprotic ionic liquids to replace harmful and flammable polar aprotic solvents such as dimethylsulfoxide, acetonitrile and dimethylformamide. As long as the synthesis routes and purification methods are more complicated, the costs of ionic liquids become even higher (Ozokwelu et al., 2017; Dupont et al., 2015).

General Synthesis Methods of Ionic Liquids

Lately, more ionic liquids have been synthesized by different methods. The reactions of the processes are commonly neutralization, quaternization and ion exchange reactions.

1. Acid-Base Neutralization

The exothermic neutralization reaction between equimolar Brönsted acid and Brönsted base that is performed either in a vessel equipped with a cooling jacket or in a flask immersed into a water bath, or an ice bath brings about the formation of ionic liquids through proton transfer, which is among the easiest synthesis routes. Following the reaction, remaining unreacted starting materials in ionic liquid can be readily removed from the formed product by washing with water, and the final product is dried in a vacuum oven (Wasserscheid and Welton, 2002). When carboxylic acids are treated with some amines at high temperatures, amides can form in place of ionic liquid.

23 more pages are available in the full version of this document, which may be purchased using the "Add to Cart" button on the publisher's webpage: www.igi-global.com/chapter/the-importance-of-ionic-liquids-and-applications-on-their-molecular-modeling/227277

Related Content

Human Factors in Dynamic E-Health Systems and Digital Libraries

Arash Shaban-Nejad and Volker Haarslev (2010). *Biomedical Knowledge Management: Infrastructures and Processes for E-Health Systems* (pp. 192-203). www.irma-international.org/chapter/human-factors-dynamic-health-systems/42607

Development of an Interactive GUI Tool for Thyroid Uptake Studies using Gamma Camera

Amruthavakkula Shiva, Vignesh T. Sai, Subramaniyan V. Siva, Kumar T. Rajamani and Sankara Sai S. Siva (2016). *International Journal of Biomedical and Clinical Engineering* (pp. 1-8). www.irma-international.org/article/development-of-an-interactive-gui-tool-for-thyroid-uptake-studies-using-gamma-camera/145162

Development of an Affordable Myoelectric Hand for Transradial Amputees

Alok Prakash and Shiru Sharma (2020). *International Journal of Biomedical and Clinical Engineering* (pp. 1-15). www.irma-international.org/article/development-of-an-affordable-myoelectric-hand-for-transradial-amputees/240742

Measurement of Cost and Economic Efficiency in Healthcare

Panagiotis Danilakis and Pericles Robolas (2006). *Handbook of Research on Informatics in Healthcare and Biomedicine* (pp. 419-424). www.irma-international.org/chapter/measurement-cost-economic-efficiency-healthcare/20608

Smart Textiles in Neonatal Monitoring: Enabling Unobtrusive Monitoring at the NICU

Fernando Seoane, Sibrecht Bouwstra, Juan Carlos Marquez, Johan Löfhede and Kaj Lindecrantz (2012). *Neonatal Monitoring Technologies: Design for Integrated Solutions* (pp. 41-55). www.irma-international.org/chapter/smart-textiles-neonatal-monitoring/65264