

Chapter 14

Seed pre-Activation Study by Means of LED Radiation

Alexey Bashilov

Moscow Aviation Institute, Russia

Mikhail Belyakov

 <https://orcid.org/0000-0002-4371-8042>

National Research University “MPEI” in Smolensk, Russia

ABSTRACT

To study the possibilities of pre-sowing seed activation, irradiation with LEDs emitting in the visible, violet, and near-ultraviolet ranges with a maximum of 405 nm was carried out. As a result of the growing experience, it was found that the height of wheat plants grown from the treated seeds significantly exceeds the control indicators except for the period of 45-55 days. To implement the flow, technology of seed activation with LEDs optoelectronic irradiation unit was developed. The advantages of the given installation are the energy efficiency and of seed treatment efficiency, due to of the optimal radiation spectrum selection and treatment doses.

INTRODUCTION

In the modern world there is a constant growth of the population and, as a consequence, the growing demand for quality food. One of the reserves of productivity growth is the use of presowing activation of seeds, including with the help of optical radiation (Kondrat'eva, Krasnolutsкая, Dukhtanova, & Obolensky, 2019, Blaszcak, Aziz, & Gryko, 2017, Hu, Li, & Jiang, 2007, Li, Ji, & Xu, 2013, Wu et al., 2013, Tsai, Huang, Chen, & Yue, 2017, Kakinoki, Kato, Ogawa, Nakao, Okai, & Katsuyama, 2013). Activation of plant seeds by led radiation is energy-efficient, environmentally safe, technological and economically justified. This study is devoted to solving the problem of choosing the optimal modes of led processing of plant seeds, including the optimal processing time, mode (continuous or pulse) and others.

Almost all known sources of radiation from discharge lamps to lasers were used for pre-sowing treatment of plant seeds (Borodin, 1996, Kondrat'eva, 2001, Loginov, 1986, Filippov, Bityuckij, &

DOI: 10.4018/978-1-5225-9420-8.ch014

Fedorov, 1997). However, an increasingly important role in lighting and irradiation techniques begin to play the light-emitting diodes (LEDs) – light sources, the generation of which occurs at the energy expense released by re-combination of carriers – electrons and holes – on the border of semiconductor materials with different character conductivity (Shubert, 2008). Particular interest as radiation sources for pre-sowing plant seeds treatment are LEDs of violet and near ultraviolet ranges (about 250-420nm).

In resistance terms to mechanical loads SD significantly exceed all other radiation sources. The service life of most modern LEDs in nominal mode exceeds 50,000 hours. this parameter is superior to all other LEDs types. Circuit SD is very simple. The advantages of LEDs are also: extremely high reliability, small size, environmental friendliness associated with the absence of mercury and other harmful substances, electrical safety (Ajzenberg, 2006).

Materials and Methods of Research

To study the possibilities of pre-sowing seeds activation, irradiation with LEDs emitting in the visible purple and near ultraviolet ranges with a maximum of 405 nm was carried out, which together with the power supply led module (Gaska, & Zhang, 2007, Bashilov, & Belyakov, 2011) (Fig 1). The optoelectronic module with sixteen LEDs creates irradiation of the working surface of 34 mW/m² at a distance of 55 cm.

During irradiation, the time of seed illumination was established and the exposure treatment dose was determined from the expression:

$$H = \int_0^{\tau} E(t)dt \quad (1)$$

where $E(t)$ – is the time dependence of irradiation in the seed treatment zone, τ -processing time.

In the simplest case, when the irradiation is constant during the exposure time, formula (1) takes the form:

Figure 1. Led matrix



20 more pages are available in the full version of this document, which may be purchased using the "Add to Cart" button on the publisher's webpage:

www.igi-global.com/chapter/seed-pre-activation-study-by-means-of-led-radiation/232100

Related Content

Is Agripreneurship a Mitigating Measure for Agricultural Issues in India ?

Dezy Kumari, Mohd Taqi, Mohd Ajmal and Aslam Khan (2022). *Driving Factors for Venture Creation and Success in Agricultural Entrepreneurship* (pp. 82-96).

www.irma-international.org/chapter/is-agripreneurship-a-mitigating-measure-for-agricultural-issues-in-india-/292969

Optimization of Green Hydrogel in Agriculture Based on Guar Gum by Response Surface Methodology

Elsayed Gamal Zaki, Shimaa Mohamed Elsaied, Faissal Aziz, Athir Mahmood Haddad, Aula Ali Alwattar and Khalid Aziz (2022). *Handbook of Research on Principles and Practices for Orchards Management* (pp. 33-50).

www.irma-international.org/chapter/optimization-of-green-hydrogel-in-agriculture-based-on-guar-gum-by-response-surface-methodology/309161

Futuristic Adoption of Blockchain Technology in the Smart Agriculture and Food Industry: Basic Concepts, Structure, Characteristics, Applicabilities

S. K. Geetha, Rajasekaran Thangaraj, J. K. Kiruthika, G. H. S. A. Devipriya and M. Naveenkumar (2023). *Contemporary Developments in Agricultural Cyber-Physical Systems* (pp. 122-148).

www.irma-international.org/chapter/futuristic-adoption-of-blockchain-technology-in-the-smart-agriculture-and-food-industry/327601

Measures of Project Worth

(2018). *Agricultural Finance and Opportunities for Investment and Expansion* (pp. 137-155).

www.irma-international.org/chapter/measures-of-project-worth/201764

Social Media Adoption by the Greek Dairy Industry

Sotiris Karetsos, Nikolaos Zafeirakopoulos, Evi Papadopoulou and Constantina Costopoulou (2023). *Influences of Social Media on Consumer Decision-Making Processes in the Food and Grocery Industry* (pp. 213-241).

www.irma-international.org/chapter/social-media-adoption-by-the-greek-dairy-industry/328380