

# Chapter 11

## Energy of Graphs

Harishchandra S. Ramane

Karnatak University, India

### ABSTRACT

*The energy of a graph  $G$  is defined as the sum of the absolute values of the eigenvalues of its adjacency matrix. The graph energy has close correlation with the total  $\pi$ -electron energy of molecules calculated with Huckel molecular orbital method in chemistry. A graph whose energy is greater than the energy of complete graph of same order is called hyperenergetic graph. A non-complete graph having energy equal to the energy of complete graph is called borderenergetic graph. Two non-cospectral graphs are said to be equienergetic graphs if they have same energy. In this chapter, the results on graph energy are reported. Various bounds for graph energy and its characterization are summarized. Construction of hyperenergetic, borderenergetic, and equienergetic graphs are reported.*

### INTRODUCTION

The energy of a graph is the sum of the absolute values of the eigenvalues of its adjacency matrix. It has a correlation with the total  $\pi$ -electron energy of a molecule in the quantum chemistry as calculated with the Huckel molecular orbital method (Gutman & Polansky, 1986).

Let  $G$  be a finite, simple, undirected graph with vertex set  $V(G)$  and edge set  $F(G)$ . The number of vertices of  $G$  is denoted by  $n$  and the number of edges of  $G$  is denoted by  $m$ . If  $V(G) = \{v_1, v_2, \dots, v_n\}$  then the adjacency matrix of  $G$  is a square matrix  $A(G) = [a_{ij}]$  of order  $n$  in which  $a_{ij} = 1$ , if the vertex  $v_i$  is adjacent to the vertex  $v_j$  and  $a_{ij} = 0$ , otherwise. The characteristic polynomial of  $A(G)$  denoted by  $\Phi(G; \lambda) = \det(\lambda I - A(G))$ , where  $I$  is an identity matrix of order  $n$ . The roots of the equation  $\Phi(G; \lambda) = 0$  are called the eigenvalues of  $G$  and they are labeled as  $\lambda_1, \lambda_2, \dots, \lambda_n$ . Their collection is called the spectrum of  $G$  denoted by  $Spec(G)$  (Cvetkovic, Doob & Sachs, 1980).

If  $\lambda_1, \lambda_2, \dots, \lambda_k$  are the distinct eigenvalues with respective multiplicities  $m_1, m_2, \dots, m_k$  then we write  $Spec(G) = \left( \begin{array}{cccc} \lambda_1 & \lambda_2 & \cdots & \lambda_k \\ m_1 & m_2 & \cdots & m_k \end{array} \right)$ . Since  $A(G)$  is a real symmetric matrix, its eigenvalues are real and can be ordered as  $\lambda_1 \geq \lambda_2 \geq \dots \geq \lambda_n$ .

DOI: 10.4018/978-1-5225-9380-5.ch011

Two non-isomorphic graphs are said to be cospectral if they have same spectra. Details about the graph spectra can be found in the book (Cvetkovic, Doob & Sachs, 1980) and for graph theoretic terminology one can refer the book (Harary, 1999).

One of the chemical applications of spectral graph theory is based on the correspondence between the graph eigenvalues and the molecular orbital energy level of  $\pi$ -electron in conjugated hydrocarbons (Gutman & Polansky, 1986).

The molecular graph of a hydrocarbon is obtained as follows: the carbon atoms are represented by the vertices and two vertices are adjacent if and only if there is a carbon-carbon bond. Hydrogen atoms are ignored.

Within the Huckel molecular orbital (HMO) method (Huckel & Quantentheoretische Beitrage zum Benzolproblem, 1931), the energy level of  $\pi$ -electron in molecules of conjugated hydrocarbons are related to the eigenvalues of a molecular graph as  $\varepsilon_i = \alpha + \beta_i$  where  $\alpha$  and  $\beta$  are empirical constants of the HMO model. The total energy of  $\pi$ -electrons denoted by  $E_\pi$  is

$$E_\pi = \sum_{i=1}^n g_i \varepsilon_i,$$

where  $g_i$  is the occupation number with energy  $\varepsilon_i$  and  $g_1 + g_2 + \dots + g_n = n$ . This yields


$$E_\pi = n\alpha + \beta \sum_{i=1}^n g_i \lambda_i \quad (1)$$

For majority of conjugated hydrocarbons  $g_i = 2$  if  $\lambda_i > 0$  and  $g_i = 0$  if  $\lambda_i < 0$ . Therefore Eq. (1) can be written as

$$E_\pi = n\alpha + 2\beta \sum_{+} \lambda_i,$$

where  $\sum_{+}$  indicates the summation over positive eigenvalues of the molecular graph.

Figure 1. Molecule and its molecular graph



28 more pages are available in the full version of this document, which may be purchased using the "Add to Cart" button on the publisher's webpage:

[www.igi-global.com/chapter/energy-of-graphs/235540](http://www.igi-global.com/chapter/energy-of-graphs/235540)

## Related Content

---

### Graph Classification of Graph Neural Networks

Gotam Singh Lalotra, Ashok Sharma, Barun Kumar Bhatti and Suresh Singh (2023). *Concepts and Techniques of Graph Neural Networks* (pp. 43-57).

[www.irma-international.org/chapter/graph-classification-of-graph-neural-networks/323821](http://www.irma-international.org/chapter/graph-classification-of-graph-neural-networks/323821)

### Deep Learning: Architectures and Applications

Kamaljit I. Lakhtaria and Darshankumar Modi (2019). *Handbook of Research on Deep Learning Innovations and Trends* (pp. 114-130).

[www.irma-international.org/chapter/deep-learning/227847](http://www.irma-international.org/chapter/deep-learning/227847)

### Optimization Algorithms

(2014). *Medical Diagnosis Using Artificial Neural Networks* (pp. 182-199).

[www.irma-international.org/chapter/optimization-algorithms/111010](http://www.irma-international.org/chapter/optimization-algorithms/111010)

### Control Signal Generator Based on Ultra-High Frequency Polynomial and Trigonometric Higher Order Neural Networks

(2021). *Emerging Capabilities and Applications of Artificial Higher Order Neural Networks* (pp. 416-454).

[www.irma-international.org/chapter/control-signal-generator-based-on-ultra-high-frequency-polynomial-and-trigonometric-higher-order-neural-networks/277686](http://www.irma-international.org/chapter/control-signal-generator-based-on-ultra-high-frequency-polynomial-and-trigonometric-higher-order-neural-networks/277686)

### A Survey on Deep Learning Techniques Used for Quality Process

Vanyashree Mardi, Naresh E. and Vijaya Kumar B. P. (2019). *Handbook of Research on Deep Learning Innovations and Trends* (pp. 131-152).

[www.irma-international.org/chapter/a-survey-on-deep-learning-techniques-used-for-quality-process/227848](http://www.irma-international.org/chapter/a-survey-on-deep-learning-techniques-used-for-quality-process/227848)