Chapter 5 Patient Portal Acceptance by the Elderly: Explained by the Elaboration Likelihood Model and Social Heuristics

Karoly Bozan

Duquesne University, USA

Kevin R. Parker Idaho State University, USA

Bill Davey https://orcid.org/0000-0002-3823-2144 *RMIT University, Melbourne, Australia*

ABSTRACT

The motivating factors that influence patient portal acceptance among the elderly are not well understood. Using the social heuristic theory, the elaboration likelihood model, and the unified theory of acceptance and use of technology, this study proposes a model that examines the persuasive mechanisms for the elderly to use patient portals. An empirical study involving 117 subjects in the United States was used to test the proposed model. Using the partial least squares method, social power, and imitate-thesuccessful social heuristics were found to significantly influence patient portal acceptance among the elderly. These findings indicate that older people invest less effort cognitively elaborating when presented with technology acceptance decisions and accept influence from their higher status peers from their network. Imitate-the-majority heuristics and central route processing were not found to be significant, implying that older people are more inclined to take advice from sources, which they find credible and invest less cognitive effort in evaluating the complex phenomena.

DOI: 10.4018/978-1-7998-1371-2.ch005

INTRODUCTION

The healthcare industry is paying greater attention to improving patient outcomes through better providerpatient communication (Tang et al., 2013; Chrischilles et al., 2014). One such tool that is seeing greater utilization by healthcare providers is the patient portal. Patient portals are secure online websites that allow patients to access their personal health information, such as communicate summaries of recent visits, medications, immunizations, allergies, and lab results, from any location with an Internet connection (Krist et al., 2012; HealthIT.gov at https://www.healthit.gov/). More sophisticated patient portals are also capable of scheduling non-urgent appointments, downloading and submitting forms, processing prescription refills, accepting payments, and providing viewable educational material (Goldzweig et al., 2013). Patient portals have become a valuable tool for addressing the problem of rapidly aging populations in most countries (Bierman, 2012).

The use of electronic health records (EHR) is also on the rise. A 2014 study (Hsiao & Hing, 2014) shows that about 80% of office-based physicians in the US use EHRs, and a majority of them (69%) are committed to participate in the meaningful use incentive payment program available in the United States (Blumenthal & Tavenner, 2010). One of the Stage 2 Core Set objectives to achieve meaningful use is "provide patients the ability to view online, download, and transmit their health information" (HealthIT. gov). This means that healthcare providers with EHR systems are mandated to provide patients with access to personal health information over the Internet using a secure online portal. Several other countries are introducing similar initiatives as they try to both improve health outcomes and strive for efficiencies in their health systems (Wickramasinghe, Davey, & Tatnall, 2013).

Numerous studies examine patient portal acceptance and use (Goldzweig et al., 2013; Ancker et al., 2011; Smith et al., 2015; Sarkar et al., 2011; Goel et al., 2011; Jung et al., 2011; Weppner et al., 2010). Patients with chronic conditions are more likely to accept and use patient portals (Goldzweig et al., 2013; Ancker et al., 2011; Millard & Fintak, 2002). Patient age is negatively correlated with portal adoption and use, especially among older patients (Goel et al., 2011; Jung, Padman, Shevchik, et al., 2011; Weppner et al., 2010). Roughly 20% of the US population will be over the age of 65 by 2030 due to longer life spans and aging baby boomers (CDC, 2013), and it has been established that doctor visits and medical spending increase during the final years of life (Hogan, Lunney, Gabel, & Lynn, 2001). In fact, some studies indicate a quarter of an individual's medical spending occurs in their final year (Lubitz & Riley, 1993). Hence, one can conclude that while older patients could benefit most from patient portals, they are the least likely group to use them.

While there exists a number of empirical studies that examine various factors to better understand consumer health information technology (CHIT) acceptance and use, they rely primarily on technology acceptance theories (Klein, 2007). The variety of antecedents in the competing models show that older patients are less likely to accept CHIT by making use of online health information because of less comfort, efficacy, and control (Smith et al., 2015; Or & Karsh, 2009). However, one of these antecedents is social influence, also called subjective norm (Ajzen, 1991; Venkatesh & Davis, 2000; Moore & Benbasat, 1991), which remains unexplored in regards to older patients. According to Carley and Kaufer (1993), elderly patients are likely to conform to the attitudes, norms, and beliefs of those around them, meaning that social influence may be a motivator for the elderly to adopt and use patient portals. It is therefore prudent to examine social influence in more detail beyond the technology acceptance theories.

22 more pages are available in the full version of this document, which may be purchased using the "Add to Cart" button on the publisher's webpage: www.igi-global.com/chapter/patient-portal-acceptance-by-the-elderly/244696

Related Content

Multiple Blind Watermarking Framework for Security and Integrity of Medical Images in E-Health Applications

Abdallah Soualmi, Adel Altiand Lamri Laouamer (2022). Research Anthology on Securing Medical Systems and Records (pp. 517-532).

www.irma-international.org/chapter/multiple-blind-watermarking-framework-for-security-and-integrity-of-medical-imagesin-e-health-applications/309015

Behavior Change through ICT Use: Experiences from Relatively Healthy Populations

Marieke W. Verheijden (2007). *Web Mobile-Based Applications for Healthcare Management (pp. 118-141).* www.irma-international.org/chapter/behavior-change-through-ict-use/31154

Proposed Framework for the Deployment of Telemedicine Centers in Rural Bangladesh

Raqibul Mostafa, Gazi Mehedi Hasan, A.M. Alomgir Kabirand Md Atiqur Rahman (2011). *International Journal of E-Health and Medical Communications (pp. 55-72).* www.irma-international.org/article/proposed-framework-deployment-telemedicine-centers/51621

Two Case Studies in Human Factors in Healthcare: The Nurse and Older Patient

Richard Pak, Nicole Fink, Margaux Priceand Dina Battisto (2010). *International Journal of Healthcare Delivery Reform Initiatives (pp. 17-38).* www.irma-international.org/article/two-case-studies-human-factors/53871

Multi-Dimensional Criteria for the Evaluation of E-Health Services

Alalwany Hamidand Alshawi Sarmad (2011). *Healthcare Delivery Reform and New Technologies: Organizational Initiatives (pp. 172-189).* www.irma-international.org/chapter/multi-dimensional-criteria-evaluation-health/50159