


Chapter 3

Evaluation of Multi-Temporal Sentinel-1 Dual Polarization SAR Data for Crop Type Classification

Thota Sivasankar

 <https://orcid.org/0000-0003-2422-8731>
NIIT University, India

Pavan Kumar Sharma

Annex Infotechnologies Pvt. Ltd., India

M. N. S. Ramya

Independent Researcher, USA

Pithani Venkatesh

Skymet Weather Services Pvt. Ltd., India

G. D. Bairagi

M.P. Council of Science and Technology, India

ABSTRACT

India is one of the highly populated countries, and its economy mainly depends on agriculture. The crop type classification is an essential requirement for ensuring food security, crop monitoring, and to understand the environmental consequences of cultivated ecosystems. This study exploits freely available multi-temporal SAR data for discriminating crop types, such as wheat, gram, and mustard, over Ashok Nagar district, Madhya Pradesh, India. Nine Sentinel-1 dual-polarized data acquired from

DOI: 10.4018/978-1-7998-5027-4.ch003

January 2018 to April 2018 in interferometric wide swath mode are used. Class separability analysis using Bhattacharyya Distance (BD) has been performed for multi-temporal VV and VH backscatter, log-ratio, and Radar Vegetation Index (RVI) to quantify the ability to distinguish temporal profiles of crops. RVI has shown the significant result in class separability analysis in comparison with other parameters. Crop type classification map has been generated using a support vector machine classifier with overall accuracy and Kappa coefficient of 96.32% and 0.95, respectively.

INTRODUCTION

According to the United Nations (United Nations, 2019), India is one of the top ten countries whose global population proliferate more than half of the projected growth between 2019 and 2050. It is also forecasted that India may be the highest populated country by 2027 based on the current growth rate. The steady increase in population will contemporaneously increase demand for food, which influence the future actions of mankind towards food security and nature conservation (FAO, 2009). Several studies, such as Yu et al. (2015) and Pingali et al. (2019) discussed the importance of agriculture and its need for a strategy in India to ensure future food security.

The spatial distribution of crop types and timely health status are significantly important for decision makers at regional, national and even global level (Brisco et al., 1998; Wu et al., 2015). In this regard, satellite Earth Observation (EO) data has become an essential data source. With the increase in EO satellites' operating in optical and microwave regions of the electromagnetic spectrum enable continuous land monitoring/observations. Particularly, free and open satellite data provided from Landsat and Sentinel missions have been extensively used for these applications (Wulder et al., 2012; Veloso et al., 2017; Kamilaris et al., 2017). Although, several studies (Thenkabail et al., 2005; Dheeravath et al., 2010) have shown the potential capabilities of optical multi-spectral data for crop classification. As the data is highly sensitive to the weather conditions, these approaches may not provide acceptable results over areas having frequent cloud cover or foggy conditions. Since radar remote sensing can provide data during almost all-weather conditions, this may be effectively used for un-interrupted crop growth monitoring than optical. In addition, its unique sensitivity towards structural, geometrical and dielectric properties of various components of the crop makes possible for crop identification and biophysical parameters retrieval, concurrently assess the health status (Patel and Srivastava, 2013; Gao et al., 2018; Sivasankar et al. 2018). Furthermore, the ability of radar signal to penetrate through vegetation and interact with underneath soil enables

16 more pages are available in the full version of this document, which may be purchased using the "Add to Cart" button on the publisher's webpage: www.igi-global.com/chapter/evaluation-of-multi-temporal-sentinel-1-dual-polarization-sar-data-for-crop-type-classification/257696

Related Content

A Framework for Understanding Adaptation by Manufacturing Industries

Saon Ray (2017). *Natural Resources Management: Concepts, Methodologies, Tools, and Applications* (pp. 302-313).

www.irma-international.org/chapter/a-framework-for-understanding-adaptation-by-manufacturing-industries/165299

Globalization, Governance, and Food Security: The Case of BRICS

Sebak K. Janaand Asim K. Karmakar (2017). *Natural Resources Management: Concepts, Methodologies, Tools, and Applications* (pp. 692-712).

www.irma-international.org/chapter/globalization-governance-and-food-security/165316

How Does Public Attention Influence Natural Gas Price?: New Evidence with Google Search Data

Xin Li, Jian Ma, Wei Shang, Shouyang Wangand Xun Zhang (2017). *Natural Resources Management: Concepts, Methodologies, Tools, and Applications* (pp. 1489-1506).

www.irma-international.org/chapter/how-does-public-attention-influence-natural-gas-price/165357

Harmonising CSR and Climate Change Mitigation and Adaptation Strategies to Build Community Adaptive Capacity in Bali's Tourism Sector

Putu Indah Rahmawati, Terry DeLacyand Min Jiang (2017). *Natural Resources Management: Concepts, Methodologies, Tools, and Applications* (pp. 1020-1040).

www.irma-international.org/chapter/harmonising-csr-and-climate-change-mitigation-and-adaptation-strategies-to-build-community-adaptive-capacity-in-balis-tourism-sector/165333

Durban Community's Experiences of Class-Based Urinary Diversion Toilets

Innocent Simphiwe Nojiyeza (2022). *Handbook of Research on Resource Management and the Struggle for Water Sustainability in Africa* (pp. 202-224).

www.irma-international.org/chapter/durban-communitys-experiences-of-class-based-urinary-diversion-toilets/295931