

Chapter 14

Antidiabetic Activity (Anti-Hyperglycemic Activity, Anti-Hyperlipidemic Activity)/ Agents From Medicinal Plants

Manish Singh Sansi

National Dairy Research Institute, India

Daraksha Iram

National Dairy Research Institute, India

Kapil Singh Narayan

National Dairy Research Institute, India

Sandeep Kumar

 <https://orcid.org/0000-0002-7576-9696>

National Dairy Research Institute, India

Om Prakash

National Dairy Research Institute, India

Dipanjan Misra

National Dairy Research Institute, India

ABSTRACT

*Diabetes mellitus (DM) is a chronic disease caused by inherited or acquired deficiency in insulin secretion and by decreased insulin secretion by the organ. Insulin deficiency causes the DM. Synthetic drugs are widely used in the treatment of diabetes, but they have some side effects. The antihyperglycemic and antihyperlipidemic effects of the plants are related to their ability to maintain pancreatic function. Medicinal plants constituents such as glycosides, alkaloids, terpenoids, and flavonoids mitigate DM. *B. ciliata* inhibits the α -glucosidase and α -amylase. Cinnamon extracts improve insulin receptor function by activating insulin receptor kinase and inhibiting insulin receptor phosphatase, which lead to an increase in insulin sensitivity. *Morinda lucida* also had the highest antioxidant activity, and it also inhibited the α -glucosidase. Many plants have also been shown to antihyperlipidemic effects. Finally, it can be concluded that medicinal plants have that ability to treat or prevent DM.*

DOI: 10.4018/978-1-6684-3546-5.ch014

INTRODUCTION

Diabetes is the metabolic disease of carbohydrates, proteins and fats, and a prolonged disease that increases when the pancreas is not able in a function to synthesize insulin, or the physique can't create perfect use of the insulin it produces (Kooti *et al.*, 2016). Insulin is the hormone which is synthesized by the pancreas that functions as a key to transport glucose from the foods into the cells to release energy for body functions. All the carbohydrate ingredients are breakdown into glucose in the blood. Insulin also helps to glucose get into the cells (Kumar *et al.*, 2011). Diabetes mellitus is not now a disease alternatively is the a group of metabolic disorders characterised by way of prolonged hyperglycemia, succeeding from failings in the secretion of insulin, increase thirst, extend ketonuria and ketonemia, urinary output are the collective signs of DM (Andrade-Cetto & Wiedenfeld, 2004). The estimated range of the diabetic patients 171 million globally in the 2000 is likely to be expand at least 366 million till the year 2030 (Shinde *et al.*, 2014). Over the long-standing high level of glucose are associated with injury to the physique and the failure of a tissues and organs. Though there are the range of procedures to decrease the patients of diabetes and its secondary complications, many desired natural formulation are there to reduce the cost.

Types of DM

The β -cell imperfections and the insulin resistance in the pancreas lead to the progress of 4 main types of DM.

Type-1 Diabetes

This is now denoted that, the (IDDM) Insulin dependent diabetes mellitus or the immune-mediated diabetes. In this case very little or no insulin produced. It is reason with the aid of autoimmune response which assaults the insulin-producing β -cells (Nazarian-Samani *et al.*, 2018). This is hardly brought about due to mutation in the HLA the chromosome locus is 6p21 generally known as IDDM1 (Ali *et al.*, 2006). Type-1 diabetes can have an effect on people at whichever age, however generally develops in the children or adults, young peoples. This type of diabetes the patients want injections of insulin daily to control the blood glucose levels (Shori, 2015).

These are most common a symptom of the type-1 diabetes includes: increase the dry mouth and thirsting; sudden weight loss; regular of urination; tiredness and Lack of the energy; continual hunger, unclear of the vision

Type-2 Diabetes

The Type-2 diabetes is the moreover known as the Non-Insulin Dependent Diabetes Mellitus (NIDDM). It is shows, that insulin resistance and low insulin excretion and debts, ~90% of situations of the type-2 diabetes. T2D generally takes region later the age of 40 years, The T2D is the most prevalent types of diabetes, and the numbers of the mostly ~90% of all the diabetes cases. Because the insulin can't work properly, and the blood glucose ranges keep rising, and releasing extra insulin (Apostolidis *et al.*, 2007). The many risk factors have been related with the type-2 diabetes and include: family history of the diabetes; overweight, harmful diets; physical indolence; increasing the age; increase blood pressure; impaired glucose tolerance (IGT); reduced nutrition during the pregnancy. The signs of type-2 diabetes are alike

16 more pages are available in the full version of this document, which may be purchased using the "Add to Cart" button on the publisher's webpage:

www.igi-global.com/chapter/antidiabetic-activity-anti-hyperglycemic-activity-anti-hyperlipidemic-activityagents-from-medicinal-plants/289485

Related Content

Functional Foods and Cardiac Health

Santosh Jain Passi (2019). *Complementary and Alternative Medicine: Breakthroughs in Research and Practice* (pp. 304-329).

www.irma-international.org/chapter/functional-foods-and-cardiac-health/211777

The Impact of Parental Substance Abuse on Children

Anu Dandona (2019). *Substance Abuse and Addiction: Breakthroughs in Research and Practice* (pp. 128-140).

www.irma-international.org/chapter/the-impact-of-parental-substance-abuse-on-children/219411

Rehabilitation Nursing Program for Elderly Functional Independence

Rogério Ferrinho Ferreira, Luís Amador Ribeiro and César João Fonseca (2020). *Handbook of Research on Health Systems and Organizations for an Aging Society* (pp. 102-114).

www.irma-international.org/chapter/rehabilitation-nursing-program-for-elderly-functional-independence/238274

Is mRNA indeed Useful in Clinical Management of Cervical Pathology?

Panagiotis Cherouveim, Antonios Athanasiou and Evangelos Paraskevidis (2021). *Handbook of Research on Oncological and Endoscopic Dilemmas in Modern Gynecological Clinical Practice* (pp. 121-128).

www.irma-international.org/chapter/is-mrna-indeed-useful-in-clinical-management-of-cervical-pathology/260080

The Uses of Cannabinoids in Medicine and Their Spectroscopic Applications

Alev Erand Sefa Çelik (2023). *Medical Cannabis and the Effects of Cannabinoids on Fighting Cancer, Multiple Sclerosis, Epilepsy, Parkinson's, and Other Neurodegenerative Diseases* (pp. 235-257).

www.irma-international.org/chapter/the-uses-of-cannabinoids-in-medicine-and-their-spectroscopic-applications/320050